

Fluctuating multicomponent lattice Boltzmann model

D. Belardinelli,^{1,*} M. Sbragaglia,^{1,†} L. Biferale,^{1,‡} M. Gross,^{2,3,§} and F. Varnik^{4,||}

¹*Department of Physics, University of Rome “Tor Vergata,” Via della Ricerca Scientifica 1, 00133 Rome, Italy*

²*Max-Planck-Institut für Intelligente Systeme, Heisenbergstraße 3, 70569 Stuttgart, Germany*

³*Institut für Theoretische Physik IV, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany*

⁴*Interdisciplinary Centre for Advanced Materials Simulation (ICAMS), Ruhr-Universität Bochum, Universitätsstrasse 150, 44780 Bochum, Germany*

(Received 21 October 2014; published 24 February 2015)

Current implementations of fluctuating lattice Boltzmann equations (FLBEs) describe single component fluids. In this paper, a model based on the continuum kinetic Boltzmann equation for describing multicomponent fluids is extended to incorporate the effects of thermal fluctuations. The thus obtained fluctuating Boltzmann equation is first linearized to apply the theory of linear fluctuations, and expressions for the noise covariances are determined by invoking the fluctuation-dissipation theorem directly at the kinetic level. Crucial for our analysis is the projection of the Boltzmann equation onto the orthonormal Hermite basis. By integrating in space and time the fluctuating Boltzmann equation with a discrete number of velocities, the FLBE is obtained for both ideal and nonideal multicomponent fluids. Numerical simulations are specialized to the case where mean-field interactions are introduced on the lattice, indicating a proper thermalization of the system.

DOI: [10.1103/PhysRevE.91.023313](https://doi.org/10.1103/PhysRevE.91.023313)

PACS number(s): 47.11.-j, 47.10.-g, 47.55.-t

I. BACKGROUND

Thermal fluctuations are important ingredients for a proper mesoscale description of a wide variety of flows in soft matter and biological physics [1–3]. Theoretically, thermally fluctuating mesoscopic flows are most conveniently dealt with within the framework of fluctuating hydrodynamics [4,5]. This approach, pioneered by Landau and Lifshitz for simple fluids, promotes the nonequilibrium fluxes to stochastic variables, thereby enabling their statistical mechanical description. Similar equations were then introduced to study the dynamics of the order parameter fluctuations in critical phenomena, as reviewed by Halperin and Hohenberg [6]. An important ingredient in this formulation is the fluctuation-dissipation theorem (FDT) relating the noise covariances to the Onsager coefficients of the fluid. At the mathematical level, this is best summarized by the fluctuating hydrodynamic equations of an isothermal binary mixture of two species with barycentric velocity \mathbf{v}_b and mass concentration C ,

$$\partial_t \rho_t + \nabla \cdot (\rho_t \mathbf{v}_b) = 0, \quad \partial_t \rho + \nabla \cdot (\rho \mathbf{v}_b) = \nabla \cdot (\mathcal{D} \nabla \mu + \Psi), \quad (1)$$

$$\begin{aligned} & \partial_t (\rho_t \mathbf{v}_b) + \nabla \cdot (\rho_t \mathbf{v}_b \mathbf{v}_b) \\ & = -\nabla P + \nabla \cdot \{ \eta [\nabla \mathbf{v}_b + (\nabla \mathbf{v}_b)^T] + \Sigma \}, \end{aligned} \quad (2)$$

where $\rho_t = \rho + \rho'$ is the total density, $\rho = \rho_t C$ the density of the first species, $\rho' = (1 - C)\rho$ the density of the second species, $P = P(\rho_t, C)$ the equation of state, and $\mu = \mu(\rho_t, C)$ the chemical potential driving diffusion of one species into the other. The capital greek letters denote stochastic diffusion and

momentum fluxes whose variance is fixed by the FDT to be (the superscript T denotes transposition),

$$\Psi = \sqrt{2\mathcal{D}}\tilde{\mathbf{W}}, \quad \Sigma = \sqrt{\eta k_B T}(\mathbf{W} + \mathbf{W}^T), \quad (3)$$

where k_B is the Boltzmann constant, T is the temperature, and \mathbf{W} and $\tilde{\mathbf{W}}$ are standard white-noise random Gaussian tensor and vector fields with uncorrelated components. The shear viscosity η [7] and the mass diffusion coefficient \mathcal{D} regulate the intensity of the viscous stresses and the diffusion fluxes [8]. The structure of the noise (3) guarantees the correctness of the equilibrium structure factors of the fluctuating fields. These can be obtained by linearizing the equations around a uniform reference state— $\rho_t = \rho_{t0} + \delta\rho_t$, $C = C_0 + \delta C$, $\mathbf{v}_b = \delta\mathbf{v}_b$, $P = P_0 + \delta P = P_0 + c_s^2[\delta\rho_t - \rho_{t0}\beta\delta C]$ —and then applying a spatial Fourier transform [9]. In our notations, β is known as the “solubility expansion” coefficient, $\rho_{t0}\beta = (\partial\rho_t/\partial C)_P$, while $c_s^2 = (\partial P/\partial\rho_t)_C$ is the squared speed of sound [10]. The results for the structure factors are found to be independent of the wave vector \mathbf{k} (here $\langle \cdot \rangle$ refers to the canonical ensemble average and the hat $\hat{\cdot}$ indicates Fourier-transformed fields) [9,11],

$$\begin{aligned} S_{C,C}(\mathbf{k}) &= \langle \delta\hat{C}(\mathbf{k})\delta\hat{C}(-\mathbf{k}') \rangle = \frac{k_B T}{\rho_{t0}\mu_C}, \\ S_{\rho_t,\rho_t}(\mathbf{k}) &= \langle \delta\hat{\rho}_t(\mathbf{k})\delta\hat{\rho}_t(-\mathbf{k}') \rangle = \rho_{t0}k_B T \left(\frac{1}{c_s^2} + \frac{\beta^2}{\mu_C} \right), \\ S_{C,\rho_t}(\mathbf{k}) &= \langle \delta\hat{C}(\mathbf{k})\delta\hat{\rho}_t(-\mathbf{k}') \rangle = \beta \frac{k_B T}{\mu_C}, \end{aligned} \quad (4)$$

where $\mu_C = (\partial\mu/\partial C)_P$. The structure factors for the total density ρ_t and concentration C can also be obtained by expanding a free-energy functional (associated with the thermodynamic description of the system) in the density fluctuations around equilibrium [12]. For square-gradient free-energy functionals, or, equivalently, if the pressure P and the chemical potential μ in Eqs. (1) and (2) depend on the gradients of the density and/or concentration fields, one

*belardinelli@roma2.infn.it

†sbragaglia@roma2.infn.it

‡biferale@roma2.infn.it

§gross@is.mpg.de

||fathollah.varnik@rub.de

obtains a Gaussian probability density in Fourier space, with a variance given by an *Ornstein-Zernike* form [1,12,13], and the structure factors acquire a dependence on \mathbf{k} [12]. Further details on the thermodynamics and fluctuating hydrodynamic equations can be found in Appendix E.

Numerical simulations of fluctuating hydrodynamic equations pose serious challenges [14–19]. Even without the presence of thermal fluctuations, modeling and simulation of multicomponent and multiphase fluid flows is extremely difficult, especially because of the problems in simulating complex diffusion processes, phase separation, and interface dynamics [20–22]. This has triggered the development of a whole range of innovative numerical methods to solve the Navier-Stokes equations, of which the lattice Boltzmann equation (LBE) [23,24] stands out due to the capability of handling boundary conditions associated with highly irregular geometries, its nearly ideal amenability to parallel computing, and the possibility to describe nonideal fluids with phase transitions and/or phase separation [25,26]. In the present paper, we are interested in formulating a *fluctuating LBE* (FLBE) for multicomponent fluids, showing how thermal noise can be successfully incorporated in the associated LBE.

The study of fluctuations in the continuous Boltzmann equation has a long history. The Boltzmann *stosszahlansatz* effectively removes fluctuations from the Boltzmann equation, giving a mean-field description of the fluid. However, fluctuations can be restored by promoting the Boltzmann equation into a Langevin equation, an idea that dates back to Kadomtsev [27], who first applied the Langevin approach to the Boltzmann equation of a dilute gas. It was shown later by Bixon and Zwanzig [28] and independently by Fox and Uhlenbeck [29,30] that this approach, in fact, leads to the well-known equations of fluctuating hydrodynamics [5] in the limit of large length and time scales. Alternatively, the fluctuating Boltzmann equation can also be derived from a master equation description of fluctuations in phase space [31,32]. Also, generalizations of the Boltzmann-Langevin equation to nonideal gases exist and have been discussed in various papers [33].

The idea of including noise in LBE is also an active research field, as witnessed by the various publications of the recent years [12–14,34–37]. The basic idea has been pioneered by Ladd [14], who suggested the introduction of noise on the nonconserved hydrodynamic modes, thus reproducing fluctuating viscous stresses in the corresponding hydrodynamic limit (small wave vectors). However, besides the hydrodynamic modes, there exist also higher-order degrees of freedom, the so-called “ghosts” [23]. The ghost sector, which is coupled to the transport sector at small length scales, acts as a sink for the thermal stress fluctuations, and thereby compromises the balance between fluctuation and dissipation. Consequently, the thermalization of the fluid remains incomplete. Adhikari *et al.* [35] were the first to recognize the necessity to include noise on all the nonphysical ghost modes, and Dünweg *et al.* [36] reformulated this approach to follow a detailed-balance condition description. In a subsequent work, Kaehler and Wagner [37] also explored the fluctuating LBE for nonvanishing mean velocities. All these implementations, however, consider ideal-gas descriptions. Only recently was there significant progress in extending FLBE to nonideal equations of state for single component fluids [12,13,38]. In

the works by Gross *et al.* [12,13], a theoretical framework for the FLBE is provided based on the theory of linear regression of fluctuations due to Onsager and Machlup [39,40]. It is the aim of the present paper to generalize such work to the case of multicomponent fluids. We are particularly interested in deriving a “discrete” kinetic model where, with a limited set of discrete kinetic velocities and a suitable implementation of stochastic terms, one can reproduce a proper thermalization of the various degrees of freedom. As a bonus, the correct behavior of fluctuating hydrodynamics for a multicomponent mixture would then be ensured in the hydrodynamic limit. A reference kinetic model of such a kind can be sketched by

$$\partial_t f_i^\sigma + \mathbf{c}_i \cdot \nabla f_i^\sigma = \sum_j \Lambda_{ij}^\sigma [f_j^{\text{eq}}(\rho^\sigma, \mathbf{v}^\sigma) - f_j^\sigma] + \Delta_i^\sigma + \Phi_i^\sigma + \xi_i^\sigma, \quad (5)$$

where $f_i^\sigma(\mathbf{r}, t)$ represents the i th Boltzmann distribution function for the species σ , i.e., the number of particles (times mass) of the σ th species at time t in a volume element $d\mathbf{r}$ around the point \mathbf{r} moving with the discrete velocity \mathbf{c}_i . The discrete Maxwellian distribution function $f_i^{\text{eq}}(\rho^\sigma, \mathbf{v}^\sigma)$ gives the equilibrium distribution for the system. Its parameters are the hydrodynamic quantities, such as mass density $\rho^\sigma(\mathbf{r}, t)$ and velocity $\mathbf{v}^\sigma(\mathbf{r}, t)$, computed from $f_i^\sigma(\mathbf{r}, t)$ as

$$\rho^\sigma = \sum_i f_i^\sigma, \quad \mathbf{v}^\sigma = \frac{1}{\rho^\sigma} \sum_i \mathbf{c}_i f_i^\sigma. \quad (6)$$

The σ th collision matrix Λ_{ij}^σ is at the core of the multiple relaxation time (MRT) implementation, allowing the independent relaxation of the “modes” of $f_i^\sigma(\mathbf{r}, t)$, defined as

$$m_a^\sigma(\mathbf{r}, t) = \sum_i T_{ai} f_i^\sigma(\mathbf{r}, t). \quad (7)$$

The T_{ai} are chosen as linearly independent and orthogonal with respect to a certain norm [35–37]. The use of a moment space representation has the advantage that the noise can be constructed such that individual masses and total momentum are explicitly conserved. We are interested in a study of nonideal effects in a mixture, for which a static nonhomogeneous equilibrium is, in general, expected. These nonideal effects are embodied in the forcing term $\Phi_i^\sigma(\mathbf{r}, t)$ on the right-hand side of Eq. (5). The noise terms $\xi_i^\sigma(\mathbf{r}, t)$ give rise to fluctuations. They are assumed to be zero-mean Gaussian random variables, uncorrelated in time and with constant variances (which can, however, be space dependent). The derivation of the precise expression of the noise covariance will be a central aspect of the present work. Due to diffusion effects (embedded in the term Δ_i^σ), which are absent in single component fluids, the momentum modes of each individual species are no longer conserved, whereas the total momentum strictly obeys momentum conservation. This naturally poses the question of how to treat the stochastic momentum fluxes so as to recover equilibration of the various modes of the probability distribution function, which would (in the hydrodynamic limit) reproduce the correlations provided by Eqs. (1)–(4). We need two major steps to accomplish this goal: *first*, a reference “continuum” kinetic model needs to be linearized around an equilibrium state, thus paving the way for the application of the theory of linear fluctuations [39–41], and the noise covariances

need to be determined by invoking the FDT directly at the kinetic level. An important input to a Boltzmann-Langevin model is provided by the equilibrium correlations of the dynamical variables, which we introduce using the ansatz [42]

$$\begin{aligned} \langle \delta f_i^\sigma(\mathbf{r}) \delta f_j^\varsigma(\mathbf{r}') \rangle &= \mu f_i^{\text{eq}}(\rho^\sigma(\mathbf{r}), \mathbf{0}) \delta_{ij} \delta(\mathbf{r} - \mathbf{r}') \delta_{\sigma\varsigma} \\ &+ f_i^{\text{eq}}(\rho^\sigma(\mathbf{r}), \mathbf{0}) f_j^{\text{eq}}(\rho^\varsigma(\mathbf{r}'), \mathbf{0}) \gamma^{\sigma\varsigma}(\mathbf{r}, \mathbf{r}'), \end{aligned} \quad (8)$$

where μ is a mass parameter and $\gamma^{\sigma\varsigma}(\mathbf{r}, \mathbf{r}')$ is the *pair correlation function* [12, 13, 42]. Equation (8) builds up on the dilute gas Poissonian property $\langle f_i^\sigma f_j^\varsigma \rangle = \delta_{ij} \delta_{\sigma\varsigma} f_i^{\text{eq}}$ and shapes the equilibrium correlations for the modes (7). *Second*, one has to properly discretize the velocity space [43–45] and control the way the FDT-based results change when moving from the “continuum” to the “discrete” model. The success of such a discrete model would naturally open the way for numerical simulations *fully* based on the LBE and compliant with the FDT.

So far, indeed, only a few studies have addressed thermal fluctuations in binary mixtures in the context of LBE. Noise-driven spinodal decomposition was studied in [19] by combining Ladd’s fluctuating LBE [14] with a fluctuating kinetic equation for the order parameter. However, this method does not ensure FDT for either the momentum or the order parameter. Thampi *et al.* [18] reported a hybrid numerical method for the solution of the model H-fluctuating hydrodynamic equations: Only the momentum conservation equations are solved using the FLBE, while finite difference and finite volumes are proposed for spatial discretization of the order parameter equation. The approach we present in this paper, instead, fully relies on LBE. Moreover, our approach is not restricted to a binary mixture but applies to a multicomponent fluid system with arbitrary number of species.

The paper is organized as follows. In Sec. II we summarize the Langevin theory of fluctuations and present key results to be used in the context of multicomponent models. In Sec. III we review the basic features of our multicomponent model in the framework of the continuum Boltzmann equation, specializing to the case of a binary mixture with two species. The kinetic model will be properly reshaped in a form that is suitable for studying the problem of fluctuations. Next, we discuss the linearization of the resulting kinetic model including MRT. The main new contribution of the present work is presented in Sec. IV, where we provide expressions for the noise covariances based on the FDT formulation of the linearized Langevin equation. In Sec. V we show how the fluctuating Boltzmann equation may be integrated in time and space to obtain the FLBE with a discrete number of velocities. When moving from the continuum model to the discrete model, special attention will be devoted to the corrections arising in both the stochastic terms and the equilibrium velocities. Numerical results and benchmarks are presented in Sec. VI, while conclusions follow in Sec. VII. Technical details and expressions for the more general case of a multicomponent system are reported in Appendixes A–D.

II. LINEAR THEORY OF FLUCTUATIONS

Crucial for our work is the theory of linear regression of fluctuations, as originally proposed by Onsager and

Machlup [39, 40]. Such theory treats fluctuations as variables which are either even or odd under time-reversal symmetry. The Boltzmann equation has a mixed character, due to the presence of the reversible advective term and the irreversible relaxation towards equilibrium (below). Fox and Uhlenbeck, therefore, generalized the Onsager and Machlup theory to such situations [29]. They consider fluctuations of a set $\{a_1, a_2, \dots, a_n\}$ of Gaussian random variables a_i with vanishing mean and probability distribution function (pdf) at equilibrium given by

$$P^{\text{eq}}\{a_1, a_2, \dots, a_n\} = \frac{1}{Z} \exp \left[-\frac{1}{2} \sum_{ij} a_i^* (G^{-1})_{ij} a_j \right], \quad (9)$$

where Z is the normalization constant of the pdf and $*$ denotes complex conjugation. The matrix of the correlations is fixed by the *entropy matrix* G_{ij} :

$$G_{ij} = \langle a_i a_j^* \rangle. \quad (10)$$

The complex dynamical variables $a_1(t), a_2(t), \dots, a_n(t)$ are then taken to obey linear Langevin equations of the form

$$\partial_t a_i(t) = - \sum_j L_{ij} a_j(t) + \xi_i(t), \quad (11)$$

where L_{ij} is a constant squared matrix having eigenvalues with a (strictly) positive real part. The $\xi_i(t)$ are assumed to be Gaussian and uncorrelated in time, with vanishing mean and equilibrium covariance,

$$\langle \xi_i(t) \xi_j(t')^* \rangle = \Xi_{ij} \delta(t - t'). \quad (12)$$

The noise matrix Ξ_{ij} is fixed by virtue of the FDT to be [46]

$$\Xi_{ij} = \sum_k (G_{ik} L_{jk}^* + L_{ik} G_{kj}). \quad (13)$$

If Eqs. (11) are integrated starting from an arbitrary initial state, the construction of the noise ensures that the proper thermal equilibrium state characterized by Eq. (10) is reached in the limit $t \rightarrow \infty$. We remark that, in the context of Langevin-type equations such as Eq. (11), the notation $\langle \cdot \rangle$ refers to the ensemble average over all possible trajectories and is equivalent to the canonical ensemble average [Eqs. (4)], provided that Eqs. (12) and (13) are satisfied. This theoretical framework provides the basis to treat fluctuations in multicomponent fluids, as already done in the case of single component fluids [12, 13]. The major contribution of this paper is to propose a linearized version of kinetic models for multicomponent systems, with both diffusion and nonideal forces included, in a form similar to (11). This paves the way to the application of Eq. (13) to predict the noise covariances in the kinetic model.

III. ISOTHERMAL MODEL FOR NONIDEAL BINARY MIXTURES

In this section we provide the essential features of the kinetic model for nonideal multicomponent fluids, focusing our attention on a binary mixture with two species. We explicitly refer to the species indices σ , representing either the first or the second species, and $\sigma' \neq \sigma$. Moreover, when possible, in the aid of a lighter and more compact notation, we refer to *unprimed* and *primed* variables instead of using

explicitly σ and σ' , respectively. The two-species model will also be directly benchmarked against numerical simulations (Sec. VI). In order to highlight nonideal effects due to thermal fluctuations, we neglect differences in molecular masses by setting each of them equal to μ . Actually, a proper generalization to different masses can be achieved by following the reference papers [47,48]. The Boltzmann distribution functions are $f(\mathbf{c}, \mathbf{r}, t)$ and $f'(\mathbf{c}, \mathbf{r}, t)$. They are defined in such a way that $f(\mathbf{c}, \mathbf{r}, t)/\mu$ and $f'(\mathbf{c}, \mathbf{r}, t)/\mu$ represent the number of particles of the respective species at time t in a volume element $d\mathbf{c} d\mathbf{r}$ around the point (\mathbf{c}, \mathbf{r}) in the one-particle phase space. Hydrodynamic quantities, such as mass densities ρ and ρ' , velocities \mathbf{v} and \mathbf{v}' , momentum densities \mathbf{j} and \mathbf{j}' , and momentum transport densities $\mathbf{\Pi}$ and $\mathbf{\Pi}'$, can then be constructed by taking suitable moments in velocity space, namely,

$$\rho = \int d\mathbf{c} f, \quad \mathbf{j} = \rho \mathbf{v} = \int d\mathbf{c} \mathbf{c} f, \quad \rho \mathbf{\Pi} = \int d\mathbf{c} \mathbf{c} \mathbf{c} f, \quad (14)$$

with similar expressions for the primed variables. Total density of mass and momentum are then given by $\rho_t = \rho + \rho'$ and $\mathbf{j}_t = \mathbf{j} + \mathbf{j}'$, respectively, while the barycentric velocity, $\mathbf{v}_b = \mathbf{j}_t/\rho_t$, is given by

$$\mathbf{v}_b = \frac{\rho \mathbf{v} + \rho' \mathbf{v}'}{\rho + \rho'} = C \mathbf{v} + C' \mathbf{v}'. \quad (15)$$

$C = \rho/\rho_t$ and $C' = \rho'/\rho_t = 1 - C$ are the concentrations of the two species. The derivation of the isothermal multicomponent model for the binary mixture starts from the evolution equation

$$\partial_t f + \mathbf{v} \cdot \nabla f = \frac{1}{\tau} [f^{\text{eq}}(\rho, \mathbf{v}_b) - f] \quad (16)$$

and similarly for the primed species (this ever-present replacement prescription is understood in what follows). Equation (16) is the celebrated single-relaxation time Bhatnagar-Gross-Krook (BGK) [49] approximation of the Boltzmann equation. Here $\tau = \tau'$ is the relaxation time characterizing the approach towards the local equilibrium $f^{\text{eq}}(\rho, \mathbf{v}_b)$, where

$$f^{\text{eq}}(\rho, \mathbf{v}; \mathbf{c}) = \frac{\rho}{(2\pi c_\tau^2)^{D/2}} \exp\left(-\frac{1}{2c_\tau^2} |\mathbf{c} - \mathbf{v}|^2\right) \quad (17)$$

is the Maxwellian distribution function. We can then identify c_τ as the ideal speed of sound at (common) temperature T :

$$c_\tau^2 = \frac{k_B T}{\mu}. \quad (18)$$

A stationary *homogeneous* equilibrium solution of Eq. (16) is $f = f^{\text{eq}}(\rho_0, \mathbf{0})$, for some constant equilibrium density ρ_0 . However, for nonideal mixtures a *nonhomogeneous* equilibrium is, in general, expected. Nonideal effects can be taken into account by adding on the right-hand side of Eq. (16) a forcing term Φ of the form

$$\Phi = -\mathbf{a} \cdot \nabla_{\mathbf{c}} f, \quad \mathbf{a} = -c_\tau^2 (\alpha_0 \nabla \rho' + \alpha_1 \Delta \nabla \rho'), \quad (19)$$

where $\Delta = \nabla \cdot \nabla$ is the Laplacian operator and \mathbf{a} is the acceleration due to a body force acting on the particles, with α_0, α_1 suitable constants. We can justify the form of the body-force acceleration based on a mean-field theory [50–54]:

The term proportional to α_0 in (19) reflects the bulk pressure of the model and controls phase separation, while the term proportional to α_1 establishes a diffuse interface whenever phase separation is achieved in the model. To highlight only nonideal effects in the mixture, we neglected all external fields and nonideal self-interactions, by assuming $\mathbf{a}(\mathbf{r}, t)$ to depend on space and time only through spatial derivatives of the mass density $\rho'(\mathbf{r}, t)$. The factor c_τ^2 is there just for later notational convenience and can always be reabsorbed by redefining the constant coefficients α_0 and α_1 . Furthermore, we assume $\alpha'_0 = \alpha_0$ and $\alpha'_1 = \alpha_1$ so that the expression of \mathbf{a}' in terms of ρ is exactly the same. Equation (16) then becomes

$$\partial_t f + \mathbf{v} \cdot \nabla f = \frac{1}{\tau} [f^{\text{eq}}(\rho, \mathbf{v}_b) - f] + \Phi. \quad (20)$$

We seek a stationary equilibrium solution of Eq. (20) of the form $f(\mathbf{r}) = f^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0})$ for some equilibrium density field $\rho_0(\mathbf{r})$ to be determined. By inserting $f = f^{\text{eq}}(\rho_0, \mathbf{0})$ in Eq. (20), we then obtain the static density profile from

$$c_\tau^2 \nabla \rho_0 = \rho_0 \mathbf{a}_0. \quad (21)$$

This equation is a condition that has to be satisfied at equilibrium by the mass density $\rho_0(\mathbf{r})$, when the body-force acceleration \mathbf{a}_0 is given by Eq. (19) computed at equilibrium. More explicitly, combining (19) and (21), we get

$$\nabla \ln \rho_0 + \alpha_0 \nabla \rho'_0 + \alpha_1 \Delta \nabla \rho'_0 = 0. \quad (22)$$

In order to *isolate* the two physical effects of *diffusive forcing* and *nonideal forcing*, we prefer to rewrite Eq. (20) as

$$\partial_t f + \mathbf{v} \cdot \nabla f = \frac{1}{\tau} [f^{\text{eq}}(\rho, \mathbf{v}) - f] + \Delta + \Phi, \quad (23)$$

where we have isolated the diffusion into the term $\Delta = \frac{1}{\tau} [f^{\text{eq}}(\rho, \mathbf{v}_b) - f^{\text{eq}}(\rho, \mathbf{v})]$, thus leaving local momentum conservation in $\frac{1}{\tau} [f^{\text{eq}}(\rho, \mathbf{v}) - f]$.

In order to apply the Langevin theory summarized in Sec. II, we have to satisfy two requirements. *First*, the evolution equation has to be linear with respect to the dynamical variables. Equation (23) is only apparently linear, the nonlinearity being hidden in $f^{\text{eq}}(\rho, \mathbf{v})$, Δ and Φ , which are nonlinear functionals of the distribution functions f and f' . The linearization is indeed discussed in Sec. IV. *Second*, the evolution equation must be an ordinary differential equation in time [Eq. (11)]. Equation (23) involves partial derivatives with respect to \mathbf{c} and \mathbf{r} , instead. These difficulties can be overcome by transforming the space gradient ∇ into a multiplicative operator in Fourier space and by working with velocity moments, as we discuss in the following. The (velocity) moments m_a ($a = 0, 1, 2, \dots$) of the Boltzmann distribution function f are defined by the relations

$$m_a(\mathbf{r}, t) = \int d\mathbf{c} T_a(\mathbf{c}) f(\mathbf{c}, \mathbf{r}, t), \quad (24)$$

$$f(\mathbf{c}, \mathbf{r}, t) = \omega(\mathbf{c}) \sum_a \frac{T_a(\mathbf{c}) m_a(\mathbf{r}, t)}{N_a},$$

where $\omega = f^{\text{eq}}(\rho, \mathbf{0})/\rho$. We refer to the T_a as *modes*. They are related to the independent components of the dimensional *Hermite polynomials*, as explained in Appendix A. The first modes are chosen in such a way that the first m_a are related

to ρ , the D component of \mathbf{v} , and the $D(D+1)/2$ independent components of $\mathbf{\Pi}$ as

$$m_0 = \rho, \quad m_\alpha = \rho v_\alpha, \quad m_{\theta_{\alpha\beta}} = \rho(\Pi_{\alpha\beta} - c_T^2 \delta_{\alpha\beta}). \quad (25)$$

The index $\theta_{\alpha\beta} = D + \min(\alpha, \beta) + |\alpha - \beta|(2D - |\alpha - \beta| + 1)/2$ goes from $D+1$ to $D(D+3)/2$ as α and β go from 1 to D . Notice that $\theta_{\alpha\beta}$ is an index of modes, while α and β are spatial indices (more details are given in Appendix A). Correspondingly, we have

$$T_0(\mathbf{c}) = 1, \quad T_\alpha(\mathbf{c}) = c_\alpha, \quad T_{\theta_{\alpha\beta}}(\mathbf{c}) = c_\alpha c_\beta - c_T^2 \delta_{\alpha\beta}. \quad (26)$$

Crucial are the orthogonality and completeness relations, respectively,

$$\int d\mathbf{c} \omega(\mathbf{c}) T_a(\mathbf{c}) T_b(\mathbf{c}) = N_a \delta_{ab}, \quad (27)$$

$$\omega(\mathbf{c}) \sum_a \frac{T_a(\mathbf{c}) T_a(\mathbf{c}')}{N_a} = \delta(\mathbf{c} - \mathbf{c}'),$$

where the N_a are normalization constants. In particular, $N_0 = 1$, $N_\alpha = c_T^2$, and $N_{\theta_{\alpha\beta}} = c_T^4(1 + \delta_{\alpha\beta})$. More explicitly, the first terms in the expansion of the Boltzmann distribution function given in Eq. (24) read

$$f = \omega \sum_a \frac{T_a m_a}{N_a}$$

$$= \omega \rho \left[1 + \frac{\mathbf{c} \cdot \mathbf{v}}{c_T^2} + \frac{(\mathbf{c}\mathbf{c} - c_T^2 \mathbb{1}) : (\mathbf{\Pi} - c_T^2 \mathbb{1})}{2c_T^4} + \dots \right], \quad (28)$$

$\mathbb{1}$ being the $D \times D$ identity. Analogously,

$$f^{\text{eq}}(\rho, \mathbf{v}) = \omega \sum_a \frac{T_a m_a^{\text{eq}}}{N_a}$$

$$= \omega \rho \left[1 + \frac{\mathbf{c} \cdot \mathbf{v}}{c_T^2} + \frac{(\mathbf{c}\mathbf{c} - c_T^2 \mathbb{1}) : \mathbf{v}\mathbf{v}}{2c_T^4} + \dots \right] \quad (29)$$

and thus

$$\Delta = \omega \sum_a \frac{T_a \Delta_a}{N_a}$$

$$= \frac{\omega \rho}{\tau} \left[1 + \frac{\mathbf{c} \cdot (\mathbf{v}_b - \mathbf{v})}{c_T^2} + \frac{(\mathbf{c}\mathbf{c} - c_T^2 \mathbb{1}) : (\mathbf{v}_b \mathbf{v}_b - \mathbf{v}\mathbf{v})}{2c_T^4} + \dots \right]. \quad (30)$$

Finally, the body-force term in (23) should also be projected onto the Hermite basis. This term involves derivatives in \mathbf{c} and cannot be expressed directly using the values of the distribution function alone. Its expansion in Hermite polynomials can be obtained from the expansion of f by taking the derivative and using (A2) [43,44]:

$$\Phi = \omega \sum_a \frac{T_a \Phi_a}{N_a} = \omega \rho \left[\frac{\mathbf{c} \cdot \mathbf{a}}{c_T^2} + \frac{(\mathbf{c}\mathbf{c} - c_T^2 \mathbb{1}) : \mathbf{a}\mathbf{v}}{c_T^4} + \dots \right]. \quad (31)$$

In what follows, we prefer not to work with formal expansions, but rather keep the forms on the right-hand side of (28)–(31) to highlight explicitly the various contributions of the relevant

modes at the level of the hydrodynamic equations (i.e., density, momentum, transport modes). The projection of the kinetic equation onto the various modes naturally paves the way for a modification of (23), by allowing independent relaxation of the modes towards equilibrium,

$$\partial_t f + \mathbf{v} \cdot \nabla f = \Lambda[f^{\text{eq}}(\rho, \mathbf{v}) - f] + \Delta + \Phi, \quad (32)$$

where Δ is now meant to be a generalized diffusive forcing given by

$$\Delta = \Lambda[f^{\text{eq}}(\rho, \mathbf{v}_b) - f^{\text{eq}}(\rho, \mathbf{v})]. \quad (33)$$

Here Λ is a linear integral operator in the velocity space, defined by

$$(\Lambda f)(\mathbf{c}) = \int d\mathbf{c}' \Lambda(\mathbf{c}, \mathbf{c}') f(\mathbf{c}'), \quad (34)$$

$$\Lambda(\mathbf{c}, \mathbf{c}') = \omega(\mathbf{c}) \sum_a \lambda_a \frac{T_a(\mathbf{c}) T_a(\mathbf{c}')}{N_a},$$

with some positive constant λ_a . In the BGK—single relaxation time—approximation we would have $\lambda_a = 1/\tau$ and thus $\Lambda(\mathbf{c}, \mathbf{c}') = \delta(\mathbf{c} - \mathbf{c}')/\tau$ [49]. Equation (32) written in terms of the moments now reads

$$\partial_t m_a + \sum_b \partial_{ab} m_b = \lambda_a [m_a^{\text{eq}}(\rho, \mathbf{v}) - m_a] + \Delta_a + \Phi_a, \quad (35)$$

where

$$\partial_{ab} = \frac{1}{N_b} \int d\mathbf{c} \omega(\mathbf{c}) T_a(\mathbf{c}) T_b(\mathbf{c}) \mathbf{v} \cdot \nabla \quad (36)$$

is a linear differential operator. Even if the single-species momentum densities $j_\alpha = m_\alpha$ and $j'_\alpha = m'_\alpha$ are not conserved, the physics requires conservation of total momentum $j_{i\alpha} = m_\alpha + m'_\alpha$ in the absence of nonideal forcing, as well as conservation of total density $\rho_t = m_0 + m'_0$. The latter is ensured by the conservation of $\rho = m_0$ and $\rho' = m'_0$ separately. The conservation of total momentum density is enforced by choosing $\lambda_\alpha = \lambda'_\alpha$. We conveniently set λ_α and λ'_α equal to some *diffusion-relaxation frequency* λ_d ,

$$\lambda_\alpha = \lambda'_\alpha = \lambda_d, \quad (\alpha = 1, \dots, D). \quad (37)$$

IV. FLUCTUATING BOLTZMANN EQUATION FOR NONIDEAL BINARY MIXTURES

We now derive a central result of the present work: the fluctuation-dissipation relation for isothermal binary mixtures. By promoting Eq. (32) into a (nonlinear) Langevin equation, we obtain

$$\partial_t f + \mathbf{v} \cdot \nabla f = \Lambda[f^{\text{eq}}(\rho, \mathbf{v}) - f] + \Delta + \Phi + \xi. \quad (38)$$

The noise term $\xi(\mathbf{c}, \mathbf{r}, t)$ gives rise to fluctuations. This is assumed to be a zero-mean Gaussian random variable, uncorrelated in time and with constant variance (which can, however, depend on \mathbf{r} and \mathbf{c}). We remark that we use for the fluctuating Boltzmann distribution function *the same* notation as for the nonfluctuating one, even if the latter is actually the ensemble average $\langle \cdot \rangle$ of the former. To avoid misunderstanding, we here denote the solution of Eq. (32) by $\langle f(\mathbf{c}, \mathbf{r}, t) \rangle$. In the equilibrium state (reached asymptotically for $t \rightarrow \infty$), the averaged distribution function reduces to the equilibrium

Maxwellian, $\langle f(\mathbf{c}, \mathbf{r}, t) \rangle \rightarrow f^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0}; \mathbf{c}) = \omega(\mathbf{c})\rho_0(\mathbf{r})$. Here, $\rho_0(\mathbf{r})$ and $\rho'_0(\mathbf{r})$ denote the average of the asymptotic mass densities, that is the solutions of Eq. (22). A useful linearization of Eq. (38) can be performed by considering perturbations around the equilibrium state at rest:

$$\rho(\mathbf{r}, t) = \rho_0(\mathbf{r}) + \delta\rho(\mathbf{r}, t), \quad \mathbf{v}(\mathbf{r}, t) = \mathbf{0} + \delta\mathbf{v}(\mathbf{r}, t). \quad (39)$$

Note that, in contrast to previous works [12,13], ρ_0 and ρ'_0 , as well as the averaged asymptotic total mass density $\rho_{r0} = \rho_0 + \rho'_0$ and concentrations $C_0 = \rho_0/\rho_{r0}$ and $C'_0 = \rho'_0/\rho_{r0} = 1 - C_0$, are now functions of the space variable \mathbf{r} , unless explicitly stated otherwise. The deviation of the Boltzmann distribution function f from its averaged asymptotic distribution $f^{\text{eq}}(\rho_0, \mathbf{0}) = \omega\rho_0$ is denoted by δf , that is,

$$\begin{aligned} \delta f(\mathbf{c}, \mathbf{r}, t) &= f(\mathbf{c}, \mathbf{r}, t) - f^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0}; \mathbf{c}) \\ &= f(\mathbf{c}, \mathbf{r}, t) - \omega(\mathbf{c})\rho_0(\mathbf{r}). \end{aligned} \quad (40)$$

By subtracting from Eq. (38) its equilibrium average, we obtain the evolution equation for δf ,

$$\partial_t \delta f + \mathbf{v} \cdot \nabla \delta f = \Lambda(\delta f^{\text{eq}} - \delta f) + \delta\Delta + \delta\Phi + \xi, \quad (41)$$

where δ indicates the deviation of a given quantity from its averaged asymptotic value (all explicit expressions are given in Appendix B). We remark that there is a difference at time t between the fluctuating deviation from equilibrium, denoted by $\delta f(\mathbf{c}, \mathbf{r}, t)$, and the fluctuation $f(\mathbf{c}, \mathbf{r}, t) - \langle f(\mathbf{c}, \mathbf{r}, t) \rangle$. These two quantities tend to coincide for large times, when the averaged system reaches equilibrium and deviations are due to fluctuations only (Fig. 1). Indeed, $\delta f(\mathbf{c}, \mathbf{r}, t)$ can be written as the sum of the fluctuating contribution $f(\mathbf{c}, \mathbf{r}, t) - \langle f(\mathbf{c}, \mathbf{r}, t) \rangle$ and a nonfluctuating deviation from equilibrium $\langle f(\mathbf{c}, \mathbf{r}, t) \rangle - f^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0}; \mathbf{c}) = \langle \delta f(\mathbf{c}, \mathbf{r}, t) \rangle$. However, in the limit of long




FIG. 1. (Color online) A schematic view of the fluctuating Boltzmann distribution $f(t)$, its ensemble average $\langle f(t) \rangle$, and its deviation from the equilibrium distribution f^{eq} . Note that, per definition, the equilibrium distribution function neither has any explicit time dependence nor exhibits any fluctuations. The ensemble averaged distribution function, $\langle f(t) \rangle$, on the other hand, is per construction free of fluctuations but may depend on time if the system is brought out of equilibrium by some perturbation. Since $\langle f(t) \rangle$ converges towards f^{eq} for long times, the difference $\delta f(t) = f(t) - f^{\text{eq}}$ becomes identical to $f(t) - \langle f(t) \rangle$ asymptotically for $t \rightarrow \infty$.

times, $\langle f(\mathbf{c}, \mathbf{r}, t) \rangle$ approaches $f^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0}; \mathbf{c})$ so that $\langle \delta f(\mathbf{c}, \mathbf{r}, t) \rangle$ approaches zero.

The linearized equation (41) can now be written in terms of the Fourier-transformed moments. To this aim, for notational convenience, we reintroduce now the indices of the species σ and σ' in such a way that $\delta\hat{m}_a^\sigma = \delta\hat{m}_a$ and $\delta\hat{m}_a^{\sigma'} = \delta\hat{m}_a'$ (the hat indicates Fourier-transformed fields). Following the derivation steps reported in Appendix B, we find a compact form,

$$\begin{aligned} \partial_t \delta\hat{m}_a^\sigma(\mathbf{k}, t) &= - \sum_{b, \zeta} \int d\mathbf{k}' \mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') \delta\hat{m}_b^\zeta(\mathbf{k}', t) + \hat{\xi}_a^\sigma(\mathbf{k}, t), \\ (\zeta &= \sigma, \sigma'), \end{aligned} \quad (42)$$

where $\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ is the integral kernel of the time-evolution matrix operator (see Appendix B for the explicit expressions). We emphasize that, despite the similarities of some of the expressions below with the multicomponent case, they concern here a binary mixture only. The general case of a mixture with more than two species is addressed in Appendixes C and D.

The evolution equation (42) is now in the form of Eq. (11), a, σ , and \mathbf{k} being the indices. The noises $\hat{\xi}_a^\sigma(\mathbf{k}, t)$ are entirely specified by the noise correlation matrix $\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$, which is defined by (for real functions of \mathbf{r} , complex conjugation is equivalent to the change $\mathbf{k} \mapsto -\mathbf{k}$ in Fourier space)

$$\langle \hat{\xi}_a^\sigma(\mathbf{k}, t) \hat{\xi}_b^\zeta(-\mathbf{k}', t') \rangle = \Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') \delta(t - t'). \quad (43)$$

For large times, the moment $\delta\hat{m}_a^\sigma(\mathbf{k}, t)$ approaches its asymptotic value $\delta\hat{m}_a^\sigma(\mathbf{k})$. Based on Eq. (13), compliance of the noise covariances with the FDT requires

$$\begin{aligned} \Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') &= \sum_{c, \kappa} \int d\mathbf{q} [\mathcal{G}_{ac}^{\sigma\kappa}(\mathbf{k}, \mathbf{q}) \mathcal{L}_{bc}^{\kappa\zeta}(-\mathbf{k}', -\mathbf{q}) \\ &\quad + \mathcal{L}_{ac}^{\sigma\kappa}(\mathbf{k}, \mathbf{q}) \mathcal{G}_{cb}^{\kappa\zeta}(\mathbf{q}, \mathbf{k}')], \end{aligned} \quad (44)$$

where the *equilibrium correlation matrix* is defined by

$$\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = \langle \delta\hat{m}_a^\sigma(\mathbf{k}) \delta\hat{m}_b^\zeta(-\mathbf{k}') \rangle. \quad (45)$$

In Appendix C we provide the following expressions for the equilibrium correlations $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$:

$$\begin{aligned} \mathcal{G}_{ab}^{\sigma\sigma}(\mathbf{k}, \mathbf{k}') &= (2\pi)^{-D/2} \mu \hat{\rho}_0(\mathbf{k} - \mathbf{k}') N_a \delta_{ab} \\ &\quad + (\hat{\rho}_0 \hat{\rho}_0 * \hat{\gamma})(\mathbf{k}, -\mathbf{k}') \delta_{a0} \delta_{b0}, \end{aligned} \quad (46)$$

$$\mathcal{G}_{ab}^{\sigma\sigma'}(\mathbf{k}, \mathbf{k}') = (\hat{\rho}_0 \hat{\rho}'_0 * \hat{\Gamma})(\mathbf{k}, -\mathbf{k}') \delta_{a0} \delta_{b0}.$$

In Eq. (46), the double asterisk $*$ denotes a normalized biconvolution in the Fourier space involving the equilibrium mass densities and the *pair correlation functions* $\gamma(\mathbf{r}, \mathbf{r}') = \gamma^{\sigma\sigma}(\mathbf{r}, \mathbf{r}')$ and $\Gamma(\mathbf{r}, \mathbf{r}') = \gamma^{\sigma\sigma'}(\mathbf{r}, \mathbf{r}')$ (Appendix C). These are related to the density structure factors $\mathcal{S}_{\rho, \rho} = \mathcal{G}_{00}^{\sigma\sigma}$ and $\mathcal{S}_{\rho, \rho'} = \mathcal{G}_{00}^{\sigma\sigma'}$ by

$$\begin{aligned} \mathcal{S}_{\rho, \rho}(\mathbf{k}, \mathbf{k}') &= \langle \delta\hat{\rho}(\mathbf{k}) \delta\hat{\rho}(-\mathbf{k}') \rangle = (2\pi)^{-D/2} \frac{k_B T}{c_T^2} \hat{\rho}_0(\mathbf{k} - \mathbf{k}') \\ &\quad + (\hat{\rho}_0 \hat{\rho}_0 * \hat{\gamma})(\mathbf{k}, -\mathbf{k}'), \end{aligned} \quad (47)$$

$$\mathcal{S}_{\rho, \rho'}(\mathbf{k}, \mathbf{k}') = \langle \delta\hat{\rho}(\mathbf{k}) \delta\hat{\rho}'(-\mathbf{k}') \rangle = (\hat{\rho}_0 \hat{\rho}'_0 * \hat{\Gamma})(\mathbf{k}, -\mathbf{k}'),$$

where Eq. (18) has been used. Momentum moments are governed by $\mathcal{S}_{j_a, j_b} = \mathcal{G}_{\alpha\beta}^{\sigma\sigma}$ and $\mathcal{S}_{j_a, j'_b} = \mathcal{G}_{\alpha\beta}^{\sigma\sigma'}$, which gives the

following results for the momentum structure factors:

$$\begin{aligned}\mathbf{S}_{\mathbf{j},\mathbf{j}}(\mathbf{k},\mathbf{k}') &= \langle \delta \hat{\mathbf{j}}(\mathbf{k}) \delta \hat{\mathbf{j}}(-\mathbf{k}') \rangle = (2\pi)^{-D/2} k_B T \hat{\rho}_0(\mathbf{k} - \mathbf{k}') \mathbb{1}, \\ \mathbf{S}_{\mathbf{j},\mathbf{j}'}(\mathbf{k},\mathbf{k}') &= \langle \delta \hat{\mathbf{j}}(\mathbf{k}) \delta \hat{\mathbf{j}}'(-\mathbf{k}') \rangle = \mathbf{0}.\end{aligned}\quad (48)$$

Furthermore, correlations between momentum and mass densities vanish. Since $\Gamma'(\mathbf{r},\mathbf{r}') = \Gamma(\mathbf{r}',\mathbf{r})$, we have three independent pair correlation functions γ , γ' , and Γ for a binary mixture. Their expressions are still unknown at this level of description. Equation (44), together with the general expression of $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k},\mathbf{k}')$ (46), constitutes the core of our results. In the following sections, we specialize these results first to the case of *homogeneous* equilibrium (Sec. IV A) and then to the case of a *nonhomogeneous* equilibrium (Sec. IV B).

A. Homogeneous equilibrium

In this case, the homogeneous condition at equilibrium $\rho_0(\mathbf{r}) = \rho_0 = \text{const}$, which clearly solves Eq. (22). As a consequence, $\hat{\rho}_0(\mathbf{k}) = (2\pi)^{D/2} \rho_0 \delta(\mathbf{k})$. All the various quantities of interest become diagonal in Fourier space (Appendix C). In particular, $\hat{\gamma}(\mathbf{k},\mathbf{k}') = (2\pi)^{D/2} \hat{\gamma}(\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}')$ and

$$\begin{aligned}\langle \xi_\alpha(\mathbf{r},t) \xi_\alpha(\mathbf{r}',t') \rangle &= 2\lambda_d k_B T \frac{\rho_0 \rho'_0}{\rho_{t0}} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (\alpha = 1, \dots, D), \\ \langle \xi_\alpha(\mathbf{r},t) \xi'_\alpha(\mathbf{r}',t') \rangle &= -2\lambda_d k_B T \frac{\rho_0 \rho'_0}{\rho_{t0}} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (\alpha = 1, \dots, D), \\ \langle \xi_a(\mathbf{r},t) \xi_a(\mathbf{r}',t') \rangle &= 2\lambda_a N_a \frac{k_B T}{c_T^2} \rho_0 \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (a = D + 1, \dots);\end{aligned}\quad (53)$$

all other noise correlations vanish. An important feature emerges from the analysis of the first two equations of (53): The noise acting on the momentum modes of primed and unprimed species are opposite, that is $\xi'_\alpha = -\xi_\alpha$ ($\alpha = 1, \dots, D$). This is just a consequence of the fact that the momentum modes of individual components *are not* conserved variables, while the *total* momentum is. The structure factors are also diagonal in Fourier space,

$$\begin{aligned}S_{\rho,\rho}(\mathbf{k}) &= \frac{k_B T}{c_T^2} \rho_0 + (2\pi)^{D/2} \rho_0^2 \hat{\gamma}(\mathbf{k}), \\ S_{\rho,\rho'}(\mathbf{k}) &= (2\pi)^{D/2} \rho_0 \rho'_0 \hat{\Gamma}(\mathbf{k}),\end{aligned}\quad (54)$$

while for the momentum fields we find

$$\begin{aligned}\mathbf{S}_{\mathbf{j},\mathbf{j}}(\mathbf{k}) &= k_B T \rho_0 \mathbb{1}, \\ \mathbf{S}_{\mathbf{j},\mathbf{j}'}(\mathbf{k}) &= \mathbf{0}.\end{aligned}\quad (55)$$

In addition, from $\rho_{t0} \delta \mathbf{v}_b = \delta \mathbf{j} + \delta \mathbf{j}'$ and Eq. (55), the structure factor for the barycentric velocity follows as

$$\mathbf{S}_{\mathbf{v}_b,\mathbf{v}_b}(\mathbf{k}) = \frac{k_B T}{\rho_{t0}} \mathbb{1}.\quad (56)$$

A few remarks regarding the pair correlation functions are also in order. The quantities $\hat{\gamma}$, $\hat{\gamma}'$, and $\hat{\Gamma}$ are usually obtained by expanding the free-energy functional (associated with the thermodynamic description of the system)

$\hat{\Gamma}(\mathbf{k},\mathbf{k}') = (2\pi)^{D/2} \hat{\Gamma}(\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}')$. Equation (45) then becomes

$$\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k},\mathbf{k}') = \langle \delta \hat{m}_a^\sigma(\mathbf{k}) \delta \hat{m}_b^\zeta(-\mathbf{k}') \rangle = G_{ab}^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}'),\quad (49)$$

with

$$\begin{aligned}G_{ab}^{\sigma\sigma}(\mathbf{k}) &= \mu \rho_0 N_a \delta_{ab} + (2\pi)^{D/2} \rho_0^2 \hat{\gamma}(\mathbf{k}) \delta_{a0} \delta_{b0}, \\ G_{ab}^{\sigma\sigma'}(\mathbf{k}) &= (2\pi)^{D/2} \rho_0 \rho'_0 \hat{\Gamma}(\mathbf{k}) \delta_{a0} \delta_{b0}.\end{aligned}\quad (50)$$

Furthermore, the expression for $\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k},\mathbf{k}')$ can be further simplified, $\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k},\mathbf{k}') = L_{ab}^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}')$ (Appendix B), and Eq. (44) reduces to

$$\Xi_{ab}^{\sigma\zeta}(\mathbf{k}) = \sum_{c,\kappa} [G_{ac}^{\sigma\kappa}(\mathbf{k}) L_{bc}^{\kappa\zeta}(-\mathbf{k}) + L_{ac}^{\sigma\kappa}(\mathbf{k}) G_{cb}^{\kappa\zeta}(\mathbf{k})],\quad (51)$$

where $\Xi_{ab}^{\sigma\zeta}(\mathbf{k})$ is defined by

$$\begin{aligned}\langle \hat{\xi}_a^\sigma(\mathbf{k},t) \hat{\xi}_b^\zeta(-\mathbf{k}',t') \rangle &= \Xi_{ab}^{\sigma\zeta}(\mathbf{k},\mathbf{k}') \delta(t - t') \\ &= \Xi_{ab}^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}') \delta(t - t').\end{aligned}\quad (52)$$

We notice now that $\Xi_{00}^{\sigma\zeta}(\mathbf{k}) = 0$. Thus, we necessarily have to set $\xi_0^\sigma(\mathbf{k},t) = 0$ identically. This makes it possible to simplify the structure of the noise covariances in real space (all calculations are reported in Appendix D 1), and the relevant noise correlations are found to be

up to the second order in the density fluctuations around homogeneous equilibrium or also by linearizing the equation of hydrodynamics [12]. By this, one obtains a Gaussian probability density in Fourier space, with a variance given by an *Ornstein-Zernike* form [1,12,13]. Instead of following such a route, we show in Appendix D 1 how the pair correlation functions can be determined by a self-consistency condition. Indeed, as noticed earlier, $\Xi_{00}^{\sigma\zeta}(\mathbf{k}) = 0$, and we necessarily have to set $\xi_0^\sigma(\mathbf{k},t) = 0$ identically. It follows that all correlations of the form $\langle \xi_0^\sigma(\mathbf{k},t) \xi_b^\zeta(-\mathbf{k}',t') \rangle$ [or equivalently $\langle \xi_a^\sigma(\mathbf{k},t) \xi_b^\zeta(-\mathbf{k}',t') \rangle$] must vanish. To be self-consistent, we then impose $\Xi_{0b}^{\sigma\zeta}(\mathbf{k}) = 0$ for any b [or equivalently $\Xi_{a0}^{\sigma\zeta}(\mathbf{k}) = 0$ for any a]. This leads to the following structure factors:

$$\begin{aligned}S_{\rho,\rho}(\mathbf{k}) &= \frac{k_B T}{c_T^2} \frac{\rho_0}{1 - \rho_0 \rho'_0 \alpha(\mathbf{k})^2}, \\ S_{\rho,\rho'}(\mathbf{k}) &= -\frac{k_B T}{c_T^2} \frac{\rho_0 \rho'_0 \alpha(\mathbf{k})}{1 - \rho_0 \rho'_0 \alpha(\mathbf{k})^2},\end{aligned}\quad (57)$$

with

$$\alpha(\mathbf{k}) = \alpha_0 - \alpha_1 |\mathbf{k}|^2.\quad (58)$$

In the absence of mutual interactions [$\alpha(\mathbf{k}) = 0$] we recover the description of two ideal gases, for which $S_{\rho,\rho}(\mathbf{k}) = k_B T \rho_0 / c_T^2$ and $S_{\rho,\rho'}(\mathbf{k}) = 0$. Instead of mass densities ρ and ρ' , we can also use the total mass density $\rho_t = \rho + \rho'$ and

concentration $C = \rho/(\rho + \rho')$ to describe the binary mixture. The associated structure factors are obtained from Eq. (57):

$$\begin{aligned} S_{\rho_i, \rho_i}(\mathbf{k}) &= \frac{\rho_{i0} k_B T}{c_T^2} \frac{1 - 2C_0(1 - C_0)\rho_{i0}\alpha(\mathbf{k})}{1 - C_0(1 - C_0)\rho_{i0}^2\alpha(\mathbf{k})^2}, \\ S_{C, C}(\mathbf{k}) &= \frac{k_B T C_0(1 - C_0)}{c_T^2 \rho_{i0}} \frac{1 + 2C_0(1 - C_0)\rho_{i0}\alpha(\mathbf{k})}{1 - C_0(1 - C_0)\rho_{i0}^2\alpha(\mathbf{k})^2}, \quad (59) \\ S_{\rho_i, C}(\mathbf{k}) &= -\frac{k_B T C_0(1 - C_0)}{c_T^2} \frac{(1 - 2C_0)\rho_{i0}\alpha(\mathbf{k})}{1 - C_0(1 - C_0)\rho_{i0}^2\alpha(\mathbf{k})^2}. \end{aligned}$$

We remark that the above results can also be obtained from the linearization of the hydrodynamic equations of motion (1) and (2). These calculations are reported in Appendix E.

$$\begin{aligned} \langle \xi_\alpha(\mathbf{r}, t) \xi_\alpha(\mathbf{r}', t') \rangle &= 2\lambda_d k_B T \frac{\rho_0(\mathbf{r}) \rho'_0(\mathbf{r})}{\rho_{i0}(\mathbf{r})} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (\alpha = 1, \dots, D), \\ \langle \xi_\alpha(\mathbf{r}, t) \xi'_\alpha(\mathbf{r}', t') \rangle &= -2\lambda_d k_B T \frac{\rho_0(\mathbf{r}) \rho'_0(\mathbf{r})}{\rho_{i0}(\mathbf{r})} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (\alpha = 1, \dots, D), \quad (60) \\ \langle \xi_a(\mathbf{r}, t) \xi_a(\mathbf{r}', t') \rangle &= 2\lambda_a N_a \frac{k_B T}{c_T^2} \rho_0(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (a = D + 1, \dots); \end{aligned}$$

all other noise correlations vanish. It is important to observe that Eqs. (60) are essentially identical to Eqs. (53) with the density fields promoted to be space-dependent variables.

V. FLUCTUATING LATTICE BOLTZMANN EQUATION

In this section we provide details for the integration of the fluctuating Boltzmann equation along the characteristics [13,55] and apply a second-order accurate scheme to evaluate the resulting integral (Sec. V A). We then discretize the velocity space (Sec. V B). These are necessary steps to promote the results discussed in the previous sections to the level of the LBE. Special attention is paid to the control of the lattice renormalizations that we have to include in the forcing and noise terms to properly use our results in the framework of the LBE. Crucial remarks are also given on the use of the noise correlations in the nonhomogeneous case [Eq. (60)].

A. Integration along characteristics

We start by writing Eq. (38) in compact notation as

$$\partial_t f + \mathbf{v} \cdot \nabla f = R, \quad (61)$$

where $R = \Lambda[f^{\text{eq}}(\rho, \mathbf{v}_b) - f] + \Phi + \xi$ is introduced for short. Note that the diffusive forcing has been reabsorbed by using the barycentric velocity \mathbf{v}_b as the argument of the Maxwellian. Given a time interval Δt , we then integrate Eq. (61) along the characteristic starting at \mathbf{r} at time t with velocity \mathbf{c} , obtaining

$$\begin{aligned} f(\mathbf{c}, \mathbf{r} + \mathbf{c}\Delta t, t + \Delta t) - f(\mathbf{c}, \mathbf{r}, t) &= \int_0^{\Delta t} ds R(\mathbf{c}, \mathbf{r} + \mathbf{c}s, t + s) \\ &\simeq \frac{\Delta t}{2} R(\mathbf{c}, \mathbf{r} + \mathbf{c}\Delta t, t + \Delta t) + \frac{\Delta t}{2} R(\mathbf{c}, \mathbf{r}, t), \quad (62) \end{aligned}$$

B. Nonhomogeneous equilibrium

At variance with the homogeneous case discussed in Sec. IV A, the background density fields $\rho_0(\mathbf{r})$ and $\rho'_0(\mathbf{r})$ are assumed now to be spatially nonhomogeneous. Based on the expression for the integral kernel $\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ in (B18), the general result for the noise covariances given in Eq. (44) can be specialized to the case of such nonhomogeneous equilibrium. The exact expression for $\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ is reported in Appendix D 2. Taking advantage of special properties of the Hermite basis functions [see relation (A19) and Appendix A], one obtains the relevant noise correlations as

where we used the trapezoidal rule to evaluate the integral. This provides an error for the evaluation of $f(\mathbf{c}, \mathbf{r} + \mathbf{c}\Delta t, t + \Delta t)$ of order $O(\Delta t^3)$. Thus, by defining the new distribution functions,

$$\bar{f} = f - \frac{\Delta t}{2} R, \quad (63)$$

and neglecting errors due to the integral evaluation, we can write

$$\bar{f}(\mathbf{c}, \mathbf{r} + \mathbf{c}\Delta t, t + \Delta t) = \bar{f}(\mathbf{c}, \mathbf{r}, t) + \Delta t R(\mathbf{c}, \mathbf{r}, t). \quad (64)$$

The next step consists of finding an expression of R in terms of \bar{f} , instead of f . This can be done by working in the moment space. From Eq. (63) we have

$$m_a = \bar{m}_a + \frac{\Delta t}{2} R_a, \quad (65)$$

where

$$R_a = \lambda_a [m_a^{\text{eq}}(\rho, \mathbf{v}_b) - m_a] + \Phi_a + \xi_a. \quad (66)$$

By inserting (65) in (66) and rearranging, we obtain

$$R_a = \bar{\lambda}_a [m_a^{\text{eq}}(\rho, \mathbf{v}_b) - \bar{m}_a] + \left(1 - \frac{\Delta t}{2} \bar{\lambda}_a\right) (\Phi_a + \xi_a), \quad (67)$$

where

$$\bar{\lambda}_a = \frac{\lambda_a}{1 + \frac{\Delta t}{2} \lambda_a}. \quad (68)$$

Based on (65) and (67), to provide closed expressions, we finally have to express $m_a^{\text{eq}}(\rho, \mathbf{v}_b)$ and Φ_a in terms of the \bar{m}_a . The equilibrium moment $m_a^{\text{eq}}(\rho, \mathbf{v}_b)$ is a function of the m_a ($a = 0, \dots, D$) only, while the body-force term Φ_a contains all the m_a . From Eq. (66), we have $R_0 = \xi_0 = 0$ and thus $\rho = \bar{\rho}$ from Eq. (65). As a bonus, from Eq. (19) the acceleration \mathbf{a} is found to be unchanged, i.e., $\mathbf{a} = \bar{\mathbf{a}}$. To compute the

Φ_a for the transport modes [$a = D + 1, \dots, D(D + 3)/2$] we need to express \mathbf{v}, \mathbf{v}' [see Eq. (31)] in terms of $\bar{\mathbf{v}}, \bar{\mathbf{v}}'$. Projecting Eq. (65) on the momentum modes ($a = 1, \dots, D$) and using the expressions (66) and (67) for R_a , gives, respectively,

$$\begin{aligned}\rho \mathbf{v} &= \rho \bar{\mathbf{v}} + \frac{\Delta t}{2} [\lambda_d \rho (\mathbf{v}_b - \mathbf{v}) + \rho \mathbf{a} + \boldsymbol{\xi}], \\ \rho \mathbf{v} &= \rho \bar{\mathbf{v}} + \frac{\Delta t}{2} \left[\bar{\lambda}_d \rho (\mathbf{v}_b - \bar{\mathbf{v}}) + \left(1 - \frac{\Delta t}{2} \bar{\lambda}_d \right) (\rho \mathbf{a} + \boldsymbol{\xi}) \right],\end{aligned}\quad (69)$$

where $(\boldsymbol{\xi})_\alpha = \xi_\alpha$, while $\bar{\lambda}_d$ and λ_d are related to by Eq. (68). By summing the first over species and using that $\boldsymbol{\xi}' = -\boldsymbol{\xi}$, we obtain

$$\rho_t \mathbf{v}_b = \rho \bar{\mathbf{v}} + \rho' \bar{\mathbf{v}}' + \frac{\Delta t}{2} (\rho \mathbf{a} + \rho' \mathbf{a}'). \quad (70)$$

As a consequence, the second becomes

$$\begin{aligned}\rho \mathbf{v} &= \rho \bar{\mathbf{v}} + \frac{\Delta t}{2} \rho \mathbf{a} - \frac{\Delta t}{2} \bar{\lambda}_d \frac{\rho \rho'}{\rho_t} \left[\bar{\mathbf{v}} - \bar{\mathbf{v}}' + \frac{\Delta t}{2} (\mathbf{a} - \mathbf{a}') \right] \\ &+ \frac{\Delta t}{2} \left(1 - \frac{\Delta t}{2} \bar{\lambda}_d \right) \boldsymbol{\xi}.\end{aligned}\quad (71)$$

Once Φ_a for the transport modes is known, one can find the expression of the transport moments m_a in terms of the

new moments \bar{m}_a by using (65)–(67). This procedure can be iterated at all orders [56]. Notice that the velocities \mathbf{v} and \mathbf{v}' , which appear in the Maxwellian $f^{\text{eq}}(\rho, \mathbf{v}_b)$ through \mathbf{v}_b , have to be properly renormalized. As we can see in Eq. (71), in principle, both noise and forcing terms play a role in this renormalization. However, Eq. (70) shows that the normalization of the barycentric velocity involves only the total body force. This is a direct consequence of the fact that $\boldsymbol{\xi} + \boldsymbol{\xi}' = \mathbf{0}$. Moreover, as emerging from Eq. (67), both the forcing and the noise need to be properly renormalized, a fact that has been already pointed out in many other papers [12–14, 35–37]. The renormalization of the forcing term is taken into account by defining

$$\bar{\Phi}_a = \left(1 - \frac{\Delta t}{2} \bar{\lambda}_a \right) \Phi_a, \quad (72)$$

where Φ_a must be expressed in terms of the new moments \bar{m}_a as explained earlier. Analogously, the renormalization of the noise takes the form

$$\bar{\xi}_a = \left(1 - \frac{\Delta t}{2} \bar{\lambda}_a \right) \xi_a. \quad (73)$$

By Eq. (60), the corresponding noise correlations become [in the homogeneous case $\rho_0(\mathbf{r}) = \rho_0 = \text{const.}$, $\rho'_0(\mathbf{r}) = \rho'_0 = \text{const.}$, and $\rho_{t0}(\mathbf{r}) = \rho_{t0} = \text{const.}$]

$$\begin{aligned}\langle \bar{\xi}_\alpha(\mathbf{r}, t) \bar{\xi}_\alpha(\mathbf{r}', t') \rangle &= (2\bar{\lambda}_d - \Delta t \bar{\lambda}_d^2) k_B T \frac{\rho_0(\mathbf{r}) \rho'_0(\mathbf{r})}{\rho_{t0}(\mathbf{r})} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (\alpha = 1, \dots, D), \\ \langle \bar{\xi}_\alpha(\mathbf{r}, t) \bar{\xi}'_\alpha(\mathbf{r}', t') \rangle &= -(2\bar{\lambda}_d - \Delta t \bar{\lambda}_d^2) k_B T \frac{\rho_0(\mathbf{r}) \rho'_0(\mathbf{r})}{\rho_{t0}(\mathbf{r})} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (\alpha = 1, \dots, D), \\ \langle \bar{\xi}_a(\mathbf{r}, t) \bar{\xi}_a(\mathbf{r}', t') \rangle &= (2\bar{\lambda}_a - \Delta t \bar{\lambda}_a^2) N_a \frac{k_B T}{c_T^2} \rho_0(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (a = D + 1, \dots),\end{aligned}\quad (74)$$

while all other noise correlations vanish. If compared with their continuum counterpart [Eq. (60)], these equations contain the extra term $-\Delta t \bar{\lambda}_a^2$ as a correction to the FDT [13, 14].

B. Discretization of the velocity space

To finally translate the results to the framework of the LBE and formulate the corresponding FLBE, we need to introduce a proper set of discrete velocities \mathbf{c}_i and the corresponding probability density function. Following a procedure that is well consolidated [43], we can write

$$f_i = \frac{w_i}{\omega(\mathbf{c}_i)} f(\mathbf{c}_i) = w_i (2\pi)^{D/2} \exp\left(-\frac{1}{2c_T^2} |\mathbf{c}_i|^2\right) c_T^D f(\mathbf{c}_i), \quad (75)$$

where the weights w_i are chosen in such a way that the integrals in velocity involving the T_a can be exactly computed from the f_i as

$$\int d\mathbf{c} T_a(\mathbf{c}) f(\mathbf{c}) = \sum_i T_a(\mathbf{c}_i) f_i. \quad (76)$$

Following [43], in order to ensure the correctness of relation (76), the dimensionless vector \mathbf{c}_i/c_T has to be equal to the i th abscissae of the Gauss-Hermite quadrature. The larger is

the number of velocities, the larger is the number of modes that we are able to reconstruct based on relation (76). We assume that a number Q of discrete velocities is large enough for the following isotropy relations to hold [43, 57]:

$$\sum_i w_i = 1, \quad \sum_i w_i \mathbf{c}_i \mathbf{c}_i = c_T^2 \mathbb{1}, \quad \sum_i w_i \mathbf{c}_i \mathbf{c}_i \mathbf{c}_i \mathbf{c}_i = 3c_T^4 \mathbb{1} \mathbb{1}. \quad (77)$$

By transforming $f(\mathbf{c}, \mathbf{r}, t)$ we obtain a set of discrete distribution functions $f_i(\mathbf{r}, t)$. Now, $f_i(\mathbf{r}, t)/\mu$ and $f'_i(\mathbf{r}, t)/\mu$ represent the number of particles of the respective species at time t in a volume $d\mathbf{r}$ around the position \mathbf{r} having velocity \mathbf{c}_i . Relations (77) are necessary and sufficient to allow the application of Eq. (76) for $a = 0, \dots, D(D + 3)/2$ to compute $\rho, \mathbf{j} = \rho \mathbf{v}$ and $\mathbf{\Pi}$ from f_i :

$$\rho = \sum_i f_i, \quad \mathbf{j} = \rho \mathbf{v} = \sum_i \mathbf{c}_i f_i, \quad \rho \mathbf{\Pi} = \sum_i \mathbf{c}_i \mathbf{c}_i f_i. \quad (78)$$

Notice that, thanks to the factor c_T^D , f_i has the same physical dimension as the mass density ρ . Let us denote with T_{ai} the discrete equivalent of $T_a(\mathbf{c}_i)$. Clearly, we have $T_{0i} = T_0(\mathbf{c}_i)$, $T_{ai} = T_a(\mathbf{c}_i)$, and $T_{\theta_{\alpha\beta}i} = T_{\theta_{\alpha\beta}}(\mathbf{c}_i)$. As a consequence of the finiteness of Q , the modes higher than transport [$a > D(D + 3)/2$], often referred to as kinetic or ghost modes, cannot be

taken as components of dimensional Hermite polynomials. This is because we want to maintain valid the orthogonality and completeness relations (27), which now become

$$\sum_i w_i T_{ai} T_{bi} = N_a \delta_{ab}, \quad w_i \sum_a \frac{T_{ai} T_{aj}}{N_a} = \delta_{ij}, \quad (79)$$

in such a way that we can define the moments m_a for $a = 0, \dots, Q - 1$ by the following invertible transformation:

$$m_a(\mathbf{r}, t) = \sum_i T_{ai} f_i(\mathbf{r}, t), \quad f_i(\mathbf{r}, t) = w_i \sum_a \frac{T_{ai} m_a(\mathbf{r}, t)}{N_a}. \quad (80)$$

The discrete transcription of Eq. (38) is the fluctuating discrete Boltzmann equation (FDBE) [12,13] for isothermal binary mixtures,

$$\partial_t f_i + \mathbf{c}_i \cdot \nabla f_i = \sum_j \Lambda_{ij} [f_j^{\text{eq}}(\rho, \mathbf{v}) - f_j] + \Delta_i + \Phi_i + \xi_i, \quad (81)$$

where the collision matrix Λ_{ij} is constructed as follows [12,13]:

$$\Lambda_{ij} = w_i \sum_a \lambda_a \frac{T_{aj} T_{ai}}{N_a}. \quad (82)$$

All the derivations of the previous sections are consequences of relations (27) and can be obtained again using (79). The discrete distribution functions $f_i(\mathbf{r}, t)$ and $f'_i(\mathbf{r}, t)$ relax for $t \rightarrow \infty$ towards fluctuating distributions equal on average to $f_i^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0}) = w_i \rho_0(\mathbf{r})$ and $f'_i(\rho'_0(\mathbf{r}), \mathbf{0}) = w_i \rho'_0(\mathbf{r})$, respectively, $\rho_0(\mathbf{r})$ and $\rho'_0(\mathbf{r})$ obeying Eq. (22).

We introduce now the dimensionless position and time variables, \mathbf{r} and n , respectively, by $\mathbf{r} = \mathbf{r} \Delta r$ and $t = n \Delta t$. Furthermore, the lattice links \mathbf{c}_i ($i = 0, \dots, Q - 1$) are defined by $\mathbf{c}_i / c_\tau = \mathbf{c}_i / c$, where

$$c = \frac{c_\tau \Delta t}{\Delta r} \quad (83)$$

is the lattice speed of sound. Once the lattice has been chosen, the Gauss-Hermite quadrature imposes a constraint on the value of c . For the D2Q9 lattice employed here ($D = 2$, $Q = 9$; see Table I) one has $c = 1/\sqrt{3}$. For notational simplicity, we will take $\Delta r = \Delta t = 1$. Thus, from the previous section, we can write the *fluctuating lattice Boltzmann equation* (FLBE) as

$$\bar{f}_i(\mathbf{r} + \mathbf{c}_i, n + 1) = \bar{f}_i(\mathbf{r}, n) + w_i \sum_a \frac{T_{ai} R_a(\mathbf{r}, n)}{N_a}, \quad (84)$$

where

$$R_a = \bar{\lambda}_a [m_a^{\text{eq}}(\rho, \bar{\mathbf{v}}_b + \frac{1}{2} \mathbf{a}_b) - \bar{m}_a] + \bar{\Phi}_a + \bar{\xi}_a, \quad (85)$$

with $\bar{\mathbf{v}}_b = \frac{\rho \bar{\mathbf{v}} + \rho' \bar{\mathbf{v}}'}{\rho + \rho'}$ and $\mathbf{a}_b = \frac{\rho \mathbf{a} + \rho' \mathbf{a}'}{\rho + \rho'}$. Furthermore,

$$\rho = \sum_i \bar{f}_i, \quad \bar{\mathbf{j}} = \rho \bar{\mathbf{v}} = \sum_i \mathbf{c}_i \bar{f}_i, \quad (86)$$

while $\bar{\Phi}_a$ and $\bar{\xi}_a$ are defined in Eqs. (72)–(74).

The use of Eq. (74) in the nonhomogeneous case (see also Sec. IV B), however, hinges on some crucial remarks. As already anticipated before, upon discretization of the velocity

TABLE I. Basis set of the D2Q9 model used in the LBE simulations. T_{ai} denotes the basis vector, N_a is the squared norm, m_a is the corresponding moment, and $\bar{\lambda}_a$ denotes its eigenvalue in the relaxation operator. The lattice speed of sound for the D2Q9 is $c = 1/\sqrt{3}$. $m_a^{\text{eq}}(\rho, \mathbf{v}) = \sum_i T_{ai} f_i^{\text{eq}}(\rho, \mathbf{v})$ is the expression for the corresponding moment of the (truncated) Maxwellian equilibrium distribution function.

a	T_{ai}	N_a	m_a	$m_a^{\text{eq}}(\rho, \mathbf{v})$	$\bar{\lambda}_a$
0	1	1	ρ	ρ	0
1	$c_{i,x}$	1/3	j_x	ρv_x	$\bar{\lambda}_j = \bar{\lambda}_d$
2	$c_{i,y}$	1/3	j_y	ρv_y	$\bar{\lambda}_j = \bar{\lambda}_d$
3	$3 c_i ^2 - 2$	4	e	$3\rho(v_x^2 + v_y^2)$	$\bar{\lambda}_e$
4	$2c_{i,x}^2 - c_i ^2$	4/9	P_{ww}	$\rho(v_x^2 - v_y^2)$	$\bar{\lambda}_s$
5	$c_{i,x} c_{i,y}$	1/9	P_{xy}	$\rho v_x v_y$	$\bar{\lambda}_s$
6	$(3 c_i ^2 - 4)c_{i,x}$	2/3	q_x	0	$\bar{\lambda}_q$
7	$(3 c_i ^2 - 4)c_{i,y}$	2/3	q_y	0	$\bar{\lambda}_q$
8	$9 c_i ^4 - 15 c_i ^2 + 2$	16	ϵ	0	$\bar{\lambda}_\epsilon$

space, one can maintain the orthogonality and completeness relations (79), but the actual form of the modes higher than transport deviates from the Hermite polynomials. A concrete example of this fact is provided by the discrete basis used in the numerical simulations of Sec. VI (Table I): One may verify explicitly that the orthogonality relations are satisfied, but the higher-order modes ($a = 6-8$) cannot be expressed as a linear combination of the Hermite polynomials of the same order, while the lower-order modes ($a = 0-5$) can. In principle, to be compliant with the theory developed, a very large set of velocities is required and the full expansion of the forcing term (31) must be considered. In practical applications this is somehow unwanted: The set of velocities is discrete and the forcing expansion (31) is usually (as we do here) truncated at the second order. Although this has no influence on the results discussed for the homogeneous equilibrium (Sec. IV A), the case of nonhomogeneous equilibrium (Sec. IV B) needs caution. Changing the structure of the Hermite polynomials as a vector basis has an effect on the structure of the noise correlations in Eq. (74), as it generates off-diagonal elements of noise between higher-order modes. In principle, these off-diagonal noise correlations have to be included in the theory to guarantee the equilibration of high-order modes. However, in order to keep the computational overhead reasonable, we prefer in the present case to perform numerical simulations based on the diagonal form of the noise given in Eqs. (74). Comparison of the so obtained results to the solutions of known problems shows generally good agreement. This is discussed in Sec. VI.

VI. NUMERICAL SIMULATIONS

Simulations of multicomponent fluids are performed using the D2Q9 lattice ($c = c_\tau = 1/\sqrt{3}$) with two species with mass densities ρ and ρ' . To perform numerical simulations, we adopt the algorithm defined by Eqs. (84) and (85). Table I shows the chosen T_{ai} and the associated modes m_a of the D2Q9 model used. The first row covers the conserved modes, i.e., the mass densities. The second and third rows cover the momentum modes. The moment e describes a bulk stress mode and the eigenvalue $\bar{\lambda}_e$ is related to the bulk viscosity. The quantities




FIG. 2. (Color online) Equilibrium ratio (ER) for the densities and the barycentric velocity as a function of wave-vector magnitude k . Simulation results are normalized according to the theoretical predictions of Eqs. (56) and (57), except for the cross-density correlation $S_{\rho,\rho'}(k)$, which is normalized by $k_B T$. The mutual interaction strength in Eq. (87) is set to $\mathcal{G} = 0$ in all the numerical simulations (ideal gases). To appreciate the effects of the noise on the momentum modes [see the first two equations in (74)], we repeated the numerical simulations by setting such noise to zero, i.e., without stochastic diffusion fluxes (no-sdf).

P_{ww} and P_{xy} are shear modes, with a common eigenvalue $\bar{\lambda}_s$ related to the shear viscosity. The ghost sector finally consists of a ghost vector current $\mathbf{q} = (q_x, q_y)$ and a ghost density mode ϵ , with eigenvalues $\bar{\lambda}_q$ and $\bar{\lambda}_\epsilon$, respectively. The body force is described on the lattice by the forces $\rho \mathbf{a}$ and $\rho' \mathbf{a}'$, with [50,51,54]

$$\mathbf{a}(\mathbf{r}) = -\mathcal{G} \sum_i w_i \rho'(\mathbf{r} + \mathbf{c}_i) \mathbf{c}_i, \quad (87)$$

where the parameter $\mathcal{G} = \mathcal{G}'$ is a coupling strength parameter regulating the intensity of the interactions. The idea of constructing forces directly on the lattice (87) is a widely used lattice formulation of an effective mean-field theory [50–52,54,58]. By Taylor expanding and using Eq. (77), we obtain the body-force-induced accelerations

$$\mathbf{a} = -c^2 \mathcal{G} \nabla \rho' - \frac{c^4 \mathcal{G}}{2} \Delta \nabla \rho' + \dots \quad (88)$$

In principle, one can neglect higher-order terms and, comparing with Eq. (19), we obtain $\alpha_0 = \alpha'_0 = \mathcal{G}$ and $\alpha_1 = \alpha'_1 = c^2 \mathcal{G}/2$. Thus, from Eq. (58) we have

$$\alpha(\mathbf{k}) = \mathcal{G} \left(1 - \frac{c^2}{2} |\mathbf{k}|^2 \right). \quad (89)$$

However, as discussed in Sec. VI A, this choice is only valid for small $|\mathbf{k}|$, while for finite $|\mathbf{k}|$ one needs to consider higher-order

terms in Eq. (88) and find a proper renormalization of $|\mathbf{k}|^2$ in (89).

With regard to the homogeneous case, we finally remark that the form of the noise for the momentum modes (74) is perfectly compatible with the stochastic fluxes of fluctuating hydrodynamics (1) and (2). As a result of the Chapman-Enskog analysis [8], we indeed recover fluctuating hydrodynamics (1) and (2), with a density-dependent diffusivity and the noise correlations (74) exactly recover Eqs. (3).

A. Equilibration ratio for homogeneous fluids

We now investigate whether the FLBE derived in the previous sections can correctly reproduce some basic statistical mechanical results in a homogeneous fluid. First, we check whether thermal noise defined by Eqs. (74) leads to the correct equilibration in a LBE simulation of a homogeneous binary mixture with resulting structure factors for the density and velocity given by (57) and (56), respectively. We test these basic results by performing simulations in a computational domain of size $L_x \times L_y = 64 \times 2$ lbu (lattice Boltzmann units) with full periodic boundary conditions. The fluctuation temperature is chosen as $T = 10^{-5}$ lbu (setting $k_B = 1$ lbu), and all the relaxation frequencies are set to $\bar{\lambda}_a = \bar{\lambda}'_a = 1$ lbu for simplicity. Uniform densities are chosen as initial condition for the simulation, $\rho = \rho' = \rho_0 = 1.0$ lbu.




FIG. 3. (Color online) Equilibrium ratio (ER) for the densities and the barycentric velocity as a function of wave-vector magnitude k . Simulation results are normalized according to the theoretical predictions of Eqs. (56) and (57), except for the cross-density correlation $S_{\rho,\rho'}(k)$, which is normalized by $k_B T$. The mutual interaction strength in Eq. (87) is set to $\mathcal{G} = 0.4$ in all the numerical simulations. For the simulation parameters chosen (see text for details), the critical point for phase separation is found at $\mathcal{G}_c = 1.0$ lbu. To appreciate the effects of the noise on the momentum modes [see the first two equations in (74)], we repeated the numerical simulations by setting such noise to zero, i.e., without stochastic diffusion fluxes (no-sdf).

The form of the noise is easily implemented in the simulations: On each lattice site we draw noise terms (independently for each mode) from a Gaussian distribution obtained from a Box-Muller algorithm [59]. Simulation results are most conveniently compared to theoretical predictions (56) and (57) by computing the equilibration ratio (ER), which is defined as the ratio of the equal-time correlations of the density and velocity divided by its expected value. This quantity is averaged over 1000 simulation snapshots. The ER is computed as a function of wave vector magnitude k along the x direction. As we are working on a lattice, it is crucial to replace the Fourier-transformed continuum Laplacian k^2 in the various equations with its discrete equivalent. The latter will be a function of k which reduces to k^2 in the limit of small k but differs from it for large wave vectors ($k \geq 1$). The discrete Fourier-transformed Laplacian can be obtained from the lattice interaction term (87), which, for the case at hand, becomes

$$a(x) = -\frac{\mathcal{G}}{6}[\rho'(x+1) - \rho'(x-1)]. \quad (90)$$

In Fourier space, the nonlocal terms produce a contribution proportional to $\sin k$. We therefore find that the term $6(1 - \sin k/k)$ plays the role of the k^2 in Eq. (89). In Fig. 2 we first investigate a situation without mutual interactions, i.e., the case of two ideal gases with mutual diffusion *only*, obtained

by setting $\mathcal{G} = 0$ in Eq. (87). To appreciate the effects of the noise on the momentum modes [$a = 1, \dots, D$, in the first two equations of (74)], we repeated the numerical simulations by setting such noise to zero, i.e., by performing the numerical simulations without stochastic diffusion fluxes (labeled as “no-sdf” in the figure). Fluctuations in the barycentric velocity are found to be independent of k and equilibrated to the theoretical value predicted by Eq. (56). A good equilibration of the velocity is found independently of the choice of the simulation scheme, i.e., with (sdf) or without (no-sdf) noise in the momentum modes (bottom right panel). However, only a proper implementation of the stochastic diffusion fluxes [Eqs. (74)] makes it possible to recover a zero cross-density correlation, $S_{\rho,\rho'}(k) = 0$ (bottom left panel) and theoretically expected self-density correlations $S_{\rho,\rho}(k)$ and $S_{\rho',\rho'}(k)$ (top panels).

In Figs. 3 and 4 we report the equilibration ratio for two cases with mutual interactions. In particular, we set $\mathcal{G} = 0.4$ (Fig. 3) and $\mathcal{G} = 0.85$ (Fig. 4). Note that the critical point at which phase separation is observed is $\mathcal{G}_c = 1$ lbu for the total background density $\rho_{t0} = 2.0$ lbu chosen [54]. Again, equilibration is found in agreement with the theoretical expectations, and the importance of the noise in the momentum modes is crucial. It is worth noting that due to the mutual interaction term, the cross-density correlation $S_{\rho,\rho'}(k)$ is different from zero.




FIG. 4. (Color online) Equilibrium ratio (ER) for the densities and the barycentric velocity as a function of wave-vector magnitude k . Simulation results are normalized according to the theoretical predictions of Eqs. (56) and (57), except for the cross-density correlation $S_{\rho,\rho'}$, which is normalized by $k_B T$. The mutual interaction strength in Eq. (87) is set to $\mathcal{G} = 0.85$ in all the numerical simulations. For the simulation parameters chosen (see text for details), the critical point for phase separation is found at $\mathcal{G}_c = 1.0$ lbu. To appreciate the effects of the noise on the momentum modes [see the first two equations in (74)], we repeated the numerical simulations by setting such noise to zero, i.e., without stochastic diffusion fluxes (no-sdf).

In Fig. 5 we report the cross-density correlation $S_{\rho,\rho'}(k)$, i.e., the diagonal part of $\langle \delta \hat{\rho}(k) \delta \hat{\rho}'(-k') \rangle$ in the homogeneous case, normalized by $k_B T$ as a function of the wave-vector magnitude k and for various interaction strength parameters \mathcal{G} . In all the cases shown, accurate agreement between our simulations and the theoretical expectations is found.

B. Capillary fluctuations in nonhomogeneous fluids

The equilibration tests previously discussed are performed in a homogeneous system. However, practical applications of multicomponent fluid simulations include phase separation where the background profile is nonhomogeneous in space. In Sec. IV B we showed that, in such a case, the local values of the densities must be used in order to compute the noises covariances defined by Eqs. (74). However, as remarked in Sec. VI, the diagonal noise correlations predicted by continuum theory [Eqs. (60)] cannot be exactly mapped onto a discrete velocity set without introducing extra off-diagonal noise terms. Nevertheless, due to the computational convenience of diagonal noise correlations, it is of interest to investigate to which extent these can be employed in nonhomogeneous situations.




FIG. 5. (Color online) The cross-density correlation $S_{\rho,\rho'}(k)$, i.e., the diagonal part of $\langle \delta \hat{\rho}(k) \delta \hat{\rho}'(-k') \rangle$ in the homogeneous case, normalized by $k_B T$ is reported as a function of the wave-vector magnitude k . The mutual interaction strength in Eq. (87) is set to $\mathcal{G} = 0.4$ (squares), $\mathcal{G} = 0.85$ (circles), $\mathcal{G} = 0.95$ (triangles). Other simulation parameters are given in the text. The critical point for phase separation is found at $\mathcal{G}_c = 1.0$ lbu. All the numerical simulations are performed with the noise on the momentum modes, i.e., with stochastic diffusion fluxes (sdf), according to Eqs. (74). The theoretical prediction of the right-hand side of Eq. (57) is also reported (red solid line). Correspondingly, we also show the prediction for the ideal-gases ($\mathcal{G} = 0$) case (blue dotted line).

A standard test case for such a fluctuating nonlinear system is represented by capillary fluctuations of a liquid-liquid interface [60,61]. Capillary fluctuations are excited by the thermal noise in the bulk and can be described (in the case of a two-dimensional problem) in terms of a local height function $h(x)$, where x denotes a position in the interfacial region [13]. In the harmonic approximation, balancing the interface energy gain due to surface tension with $k_B T$, we obtain the static spectrum of the local height fluctuations h of a flat interface,

$$\langle |\hat{h}(k)|^2 \rangle = \frac{k_B T}{\tilde{\gamma} k^2}, \quad (91)$$

where $\tilde{\gamma}$ is the surface tension and k is just the wave vector in the interfacial region. In order to test whether the static spectrum (91) can be reproduced by our fluctuating nonideal fluid model, we perform simulations of a liquid stripe in a rectangular box of size $L_x \times L_y = 100 \times 512$ lbu with full periodic boundary conditions. The extension of a stripe is taken as 50×512 lbu. The fluctuation temperature is chosen as $T = 10^{-5}$ lbu (setting $k_B = 1$ lbu), and all the relaxation frequencies are set to $\bar{\lambda}_a = \bar{\lambda}'_a = 1$ lbu for simplicity. The interaction strength parameter \mathcal{G} is chosen to be $\mathcal{G} = 1.5$ lbu, which provides phase separation with background bulk densities $\rho_0 = 2.3$ lbu and $\rho'_0 = 0.06$ lbu in the stripe region. The nonideal interface width is approximately 6 lbu. The surface tension $\tilde{\gamma}$ is independently estimated from a Laplace experiment to be $\tilde{\gamma} = 0.174$ lbu. The capillary spectrum is obtained by averaging over 2000 snapshots in a simulation running for 2×10^6 time steps. In Fig. 6, we report the static spectrum compared with the theoretical prediction given in Eq. (91): The agreement between the numerics and the theory is very good for practically all wave vectors up to $k \approx 1$. This fact suggests that, at least for the presently studied inhomogeneous situation, possible off-diagonal noise correlations emerging in the transition from the continuous to the discrete Boltzmann equation are not relevant for equilibration.




FIG. 6. (Color online) Capillary fluctuations of a planar one-dimensional interface obtained with mean-field lattice interactions (see Sec. VIB). The equal-time spectrum of interfacial height fluctuations obtained from LBE simulations (circles) is compared with the theoretical structure factor (solid line) reported in Eq. (91). The wave vector in the interface region is indicated with k . Simulation parameters are reported in the text.

VII. CONCLUSIONS

In the present paper, we propose a generalization of the work by Gross *et al.* [12,13], describing implementations of FLBE for nonideal single component fluids to the case of multicomponent fluids. Due to diffusion effects, which are absent in single component fluids, the momentum modes of each species *are not* conserved variables, while the *total* momentum is. The theoretical basis of the proposed FLBE formulation is a kinetic model for nonideal mixtures which has been promoted to a MRT model [47], by allowing the process of relaxation towards equilibrium to be controlled separately for the different modes. Technically, we first analyze the continuum Boltzmann equation, by promoting it to a linear Langevin equation, where fluctuations can be analyzed within the theory of linearized nonequilibrium thermodynamics due to Onsager and Machlup [39,40]. The noise covariances hinge on the knowledge of the linearized relaxation, diffusion and forcing, and the structure factors, the latter obtained from self-consistency arguments in the theory. The equilibrium correlations are determined by invoking results of the kinetic theory of fluids [42]. By suitable integration in time and space, we also study the corresponding LBE, with special attention to the corrections arising for the noises and body-force terms. By tuning the strength of the nonideal interactions, we investigate both the cases of *homogeneous* (Sec. IV A) and *nonhomogeneous* (Sec. IV B) equilibrium. Numerical simulations indicate a proper thermalization of the system at all the length scales investigated.

In the nonhomogeneous case, using the continuum theory (Sec. IV B), we predict the same form of the noise covariances obtained for homogeneous systems, but with the density fields promoted to space-dependent variables. This is what would have been intuitively expected based on the notion of local equilibrium. However, while noise correlations are found to be diagonal within the continuum Boltzmann description, extra off-diagonal noise terms will appear upon projection to the lattice Boltzmann level. Indeed, crucial for our analysis is the projection of the continuum Boltzmann equation onto the orthonormal Hermite basis: Upon discretization of the velocity space, one can maintain the orthogonality and completeness relations of the modes, but the actual form of the modes higher than transport deviates from the Hermite polynomials. A systematic study of the degree of equilibration of all the modes in the presence of a nonhomogeneous background density profile will be performed in a future study. This will also give an idea to which extent extra off-diagonal noise correlations are relevant for achieving thermal equilibration of the modes. It is, however, reassuring that numerical investigations of capillary fluctuations upon neglecting off-diagonal terms (Sec. VIB) reveal a good thermalization of the interface fluctuations.

While, formally, the expressions for the noise in the nonhomogeneous case are derived for a nonfluctuating background density, in practice, the actual local value of the densities have to be used in the simulations. This naturally raises the question of the proper stochastic scheme used to integrate the discrete Langevin equations (e.g., Ito or Stratonovich) [62] as well as subtle issues related to spatially dependent friction terms [63]. Such aspects have so far not been discussed in the context

of the LBE. In order to assess the FDT in the presence of nonlinearities in the forcing (such as those responsible for phase separation), it might also be interesting to adopt a Fokker-Planck treatment to the nonideal LBE (cf. [64]). These issues are reserved for future works. We finally remark that the results presented in this paper are quite general, although the simulation results have been provided only for a case of two species with mutual interactions, where mean-field forces are introduced on the lattice [50,51,54]. Changing the form of the forcing would affect the form of $\alpha(\mathbf{k})$ in Eq. (58), but most of our treatment is still applicable. Extending the simulation results to other kinds of forces [65,66] and comparing with alternative approaches [18,36] is therefore surely warranted for future investigations.

ACKNOWLEDGMENTS

M.S., D.B., and L.B. kindly acknowledge funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement No. 279004. M.S. acknowledges Professor C. Colosqui for useful discussions and exchange of ideas on fluctuating hydrodynamics during his visit in May 2014. ICAMS acknowledges funding from its industrial sponsors, the state of North-Rhine Westphalia, and the European Commission in the framework of the European Regional Development Fund (ERDF).

APPENDIX A: DIMENSIONAL HERMITE POLYNOMIALS

The n th-dimensional Hermite polynomial $H_{\alpha_1, \dots, \alpha_n}$ can be defined using the Maxwellian computed at equilibrium,

$$H_{\alpha_1, \dots, \alpha_n}(\mathbf{c} - \mathbf{v}_0) = \frac{(-c_T^2)^n}{f^{\text{eq}}(\rho_0, \mathbf{v}_0; \mathbf{c})} \partial_{c_{\alpha_1}} \cdots \partial_{c_{\alpha_n}} f^{\text{eq}}(\rho_0, \mathbf{v}_0; \mathbf{c}), \quad (\text{A1})$$

for some common hydrodynamic velocity \mathbf{v}_0 [37]. With zero velocity, $\mathbf{v}_0 = \mathbf{0}$, we obtain

$$H_{\alpha_1, \dots, \alpha_n}(\mathbf{c}) = \frac{(-c_T^2)^n}{\omega(\mathbf{c})} \partial_{c_{\alpha_1}} \cdots \partial_{c_{\alpha_n}} \omega(\mathbf{c}). \quad (\text{A2})$$

$H_{\alpha_1, \dots, \alpha_n}(\mathbf{c})$ is a polynomial in \mathbf{c} of order n and a fully symmetric tensor of rank n , with $n = 0, 1, 2, \dots$ (for $n = 0$ we impose that $\alpha_1, \dots, \alpha_0 \equiv 0$). The first few are

$$H_0(\mathbf{c}) = 1, \quad H_\alpha(\mathbf{c}) = c_\alpha, \quad H_{\alpha\beta}(\mathbf{c}) = c_\alpha c_\beta - c_T^2 \delta_{\alpha\beta}. \quad (\text{A3})$$

A recurrence relation holds [43,45]

$$c_\beta H_{\alpha_1, \dots, \alpha_n}(\mathbf{c}) = H_{\alpha_1, \dots, \alpha_n \beta}(\mathbf{c}) + c_T^2 \sum_{k=1}^n \delta_{\alpha_k \beta} H_{\alpha_1, \dots, \alpha_{k-1} \alpha_{k+1}, \dots, \alpha_n}(\mathbf{c}). \quad (\text{A4})$$

Furthermore, the following orthogonality and completeness relations hold

$$\int d\mathbf{c} \omega(\mathbf{c}) H_{\alpha_1, \dots, \alpha_n}(\mathbf{c}) H_{\mu_1, \dots, \mu_n}(\mathbf{c}) = \delta_{mn} c_T^{2n} \delta_{\alpha_1, \dots, \alpha_n, \mu_1, \dots, \mu_n}^{(n)}, \quad (\text{A5})$$

$$\omega(\mathbf{c}) \sum_{n=0}^{\infty} \sum_{\alpha_1, \dots, \alpha_n} \frac{H_{\alpha_1, \dots, \alpha_n}(\mathbf{c}) H_{\alpha_1, \dots, \alpha_n}(\mathbf{c}')}{n! c_T^{2n}} = \delta(\mathbf{c} - \mathbf{c}'), \quad (\text{A6})$$

where $\delta_{\alpha_1, \dots, \alpha_n, \mu_1, \dots, \mu_n}^{(n)}$ vanishes unless (μ_1, \dots, μ_n) is a permutation of $(\alpha_1, \dots, \alpha_n)$. In general, notice that $\delta_{\alpha_1, \dots, \alpha_n, \mu_1, \dots, \mu_n}^{(n)}$ is not equal to unity. Indeed, from (19) and (8) of [45], one gets $\delta_{\alpha_1, \dots, \alpha_n, \mu_1, \dots, \mu_n}^{(n)} = \partial_{c_{\mu_1}} \cdots \partial_{c_{\mu_n}} c_{\alpha_1} \cdots c_{\alpha_n}$. The first few are

$$\delta_0^{(0)} = 1, \quad \delta_{\alpha, \mu}^{(1)} = \delta_{\alpha\mu}, \quad \delta_{\alpha\beta, \mu\nu}^{(2)} = \delta_{\alpha\mu} \delta_{\beta\nu} + \delta_{\alpha\nu} \delta_{\beta\mu}. \quad (\text{A7})$$

Because of their full symmetry, the number of independent $H_{\alpha_1, \dots, \alpha_n}$, for a given n , is, in general, lesser than D^n . For example, for $n = 2$, the number of independent $H_{\alpha\beta}$ is $D(D+1)/2$. The modes T_a ($a = 0, 1, 2, \dots$) are defined from the $H_{\alpha_1, \dots, \alpha_n}$ in such a way that two polynomials that differ only by a permutation of distinct indices are counted one time instead of two. One can also organize the indices of the modes according to the physical meaning, i.e., if they are related to density, momentum, transport modes, etc. In this way, they become

$$a = 0, \quad a = 1, \dots, D, \quad a = D + 1, \dots, D(D + 3)/2, \dots, \quad (\text{A8})$$

where $D(D+3)/2 = D + D(D+1)/2$. Correspondingly, each subset can be identified by $\Theta^{(n)}$, that is,

$$\Theta^{(0)} = \{0\}, \quad \Theta^{(1)} = \{1, \dots, D\}, \quad \Theta^{(2)} = \{D + 1, \dots, D(D + 3)/2\}, \dots \quad (\text{A9})$$

The number of indices of modes in $\Theta^{(n)}$ is equal to the number of independent polynomials of order n . Furthermore, we define a fully symmetric tensor $\theta_{\alpha_1, \dots, \alpha_n}$ in such a way that it varies in $\Theta^{(n)}$ as $\alpha_1, \dots, \alpha_n$ go from 1 to D . For example, we can set

$$\theta_0 = 0, \quad \theta_\alpha = \alpha, \quad \theta_{\alpha\beta} = D + \min(\alpha, \beta) + \frac{1}{2}|\alpha - \beta|(2D - |\alpha - \beta| + 1), \dots \quad (\text{A10})$$

The modes T_a ($a \in \Theta^{(0)} \cup \Theta^{(1)} \cup \Theta^{(2)} \cup \dots$) are then defined by

$$T_{\theta_{\alpha_1, \dots, \alpha_n}} = H_{\alpha_1, \dots, \alpha_n}. \quad (\text{A11})$$

This leads to the definition of the moments m_a as in Eq. (24). Furthermore, orthogonality and completeness relations (A5) and (A6) can now be written in the form (27), with the squared norms N_a defined by

$$N_{\theta_{\alpha_1, \dots, \alpha_n}} = c_T^{2n} \delta_{\alpha_1, \dots, \alpha_n, \alpha_1, \dots, \alpha_n}. \quad (\text{A12})$$

The first few are

$$N_0 = 1, \quad N_\alpha = c_T^2, \quad N_{\alpha\beta} = c_T^4 (1 + \delta_{\alpha\beta}). \quad (\text{A13})$$

The constants N_a take into account the number of permutations of distinct spatial indices in the definition of the corresponding T_a . We have

$$\sum_a \frac{T_a(\mathbf{c})T_a(\mathbf{c}')}{N_a} = \sum_{n=0}^{\infty} \sum_{a \in \Theta^{(n)}} \frac{n!c_T^{2n}}{N_a} \frac{T_a(\mathbf{c})T_a(\mathbf{c}')}{n!c_T^{2n}} = \sum_{n=0}^{\infty} \sum_{\alpha_1, \dots, \alpha_n} \frac{T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c})T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c}')}{n!c_T^{2n}}. \quad (\text{A14})$$

Based on (A12)–(A14), it follows that $n!c_T^{2n}/N_a$, with $a \in \Theta^{(n)}$, is equal to the number of multi-indices $(\alpha_1, \dots, \alpha_n)$ such that $\theta_{\alpha_1, \dots, \alpha_n} = a$. The recurrence relation (A4) becomes

$$c_\beta T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c}) = T_{\theta_{\alpha_1, \dots, \alpha_n \beta}}(\mathbf{c}) + c_T^2 \sum_{k=1}^n \delta_{\alpha_k \beta} T_{\theta_{\alpha_1, \dots, \alpha_{k-1} \alpha_{k+1}, \dots, \alpha_n}}(\mathbf{c}). \quad (\text{A15})$$

Equation (A15) can be used to prove a useful relation that will be used later. Multiplying by $\omega(\mathbf{c})T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c}')/n!c_T^{2n}$ both sides and summing over $\alpha_1, \dots, \alpha_n$ and n , we obtain

$$c_\beta \delta(\mathbf{c} - \mathbf{c}') = \omega(\mathbf{c}) \sum_{n=0}^{\infty} \sum_{\alpha_1, \dots, \alpha_n} \frac{T_{\theta_{\alpha_1, \dots, \alpha_n \beta}}(\mathbf{c})T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c}') + T_{\theta_{\alpha_1, \dots, \alpha_n \beta}}(\mathbf{c}')T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c})}{n!c_T^{2n}}, \quad (\text{A16})$$

where we used that $T_{\theta_{\alpha_1, \dots, \alpha_n}}(\mathbf{c})$ is fully symmetric to replace $\sum_{k=1}^n \delta_{\alpha_k \beta} T_{\theta_{\alpha_1, \dots, \alpha_{k-1} \alpha_{k+1}, \dots, \alpha_n}}(\mathbf{c})$ with $n\delta_{\alpha_n \beta} T_{\theta_{\alpha_1, \dots, \alpha_{n-1}}}(\mathbf{c})$ when summing over all $\alpha_1, \dots, \alpha_n$. Multiplying by $\omega(\mathbf{c}')T_a(\mathbf{c})T_b(\mathbf{c}')$ both sides and integrating over $d\mathbf{c}d\mathbf{c}'$, we obtain

$$\begin{aligned} \int d\mathbf{c} \omega(\mathbf{c})T_a(\mathbf{c})T_b(\mathbf{c})c_\beta &= \sum_{n=0}^{\infty} \sum_{\alpha_1, \dots, \alpha_n} \frac{N_a \delta_{a\theta_{\alpha_1, \dots, \alpha_n \beta}} N_b \delta_{b\theta_{\alpha_1, \dots, \alpha_n}} + N_b \delta_{b\theta_{\alpha_1, \dots, \alpha_n \beta}} N_a \delta_{a\theta_{\alpha_1, \dots, \alpha_n}}}{n!c_T^{2n}} \\ &= c_T^2 (\delta_{a\beta} \delta_{b0} + \delta_{b\beta} \delta_{a0}) + c_T^4 \sum_{\alpha} [(1 + \delta_{\alpha\beta}) \delta_{a\theta_{\alpha\beta}} \delta_{b\alpha} + (1 + \delta_{\alpha\beta}) \delta_{b\theta_{\alpha\beta}} \delta_{a\alpha}] + \dots \end{aligned} \quad (\text{A17})$$

Multiplying by $V_\beta = (\mathbf{V})_\beta$ and summing over β , we then obtain

$$\mathbf{V} \cdot \int d\mathbf{c} \omega(\mathbf{c})T_a(\mathbf{c})T_b(\mathbf{c})\mathbf{c} = c_T^2 [\boldsymbol{\delta}_a \cdot \delta_{b0} \mathbf{V} + \boldsymbol{\delta}_b \cdot \delta_{a0} \mathbf{V} + c_T^2 (\boldsymbol{\tau}_a : \boldsymbol{\delta}_b \mathbf{V} + \boldsymbol{\tau}_b : \boldsymbol{\delta}_a \mathbf{V}) + \dots]. \quad (\text{A18})$$

This relation is valid for any vector \mathbf{V} . In particular, by choosing $\mathbf{V} = i\mathbf{k}$ we obtain

$$N_b A_{ab}(\mathbf{k}) = i c_T^2 [\boldsymbol{\delta}_a \cdot \delta_{b0} \mathbf{k} + \boldsymbol{\delta}_b \cdot \delta_{a0} \mathbf{k} + c_T^2 (\boldsymbol{\tau}_a : \boldsymbol{\delta}_b \mathbf{k} + \boldsymbol{\tau}_b : \boldsymbol{\delta}_a \mathbf{k}) + \dots], \quad (\text{A19})$$

$A_{ab}(\mathbf{k})$ being defined in Eq. (B13). We use Eq. (A19) in Appendix D.2.

APPENDIX B: LINEARIZATION OF THE FLUCTUATING KINETIC MODEL FOR NONIDEAL BINARY MIXTURES

We start by Eq. (41), that is,

$$\partial_t \delta f + \mathbf{v} \cdot \nabla \delta f = \Lambda(\delta f^{\text{eq}} - \delta f) + \delta \Delta + \delta \Phi + \xi, \quad (\text{B1})$$

with

$$\delta f^{\text{eq}} = \omega \left(\delta \rho + \rho_0 \frac{\mathbf{c} \cdot \mathbf{v}}{c_T^2} \right), \quad (\text{B2})$$

$$\delta \Delta = \lambda_d \omega \rho_0 \frac{\mathbf{c} \cdot (\mathbf{v}_b - \mathbf{v})}{c_T^2}, \quad (\text{B3})$$

$$\delta \Phi = \omega \left[\rho_0 \frac{\mathbf{c} \cdot \boldsymbol{\delta} \mathbf{a}}{c_T^2} + \delta \rho \frac{\mathbf{c} \cdot \mathbf{a}_0}{c_T^2} + \rho_0 \frac{(\mathbf{c}\mathbf{c} - c_T^2 \mathbf{1}) : \mathbf{a}_0 \mathbf{v}}{c_T^4} + \dots \right]. \quad (\text{B4})$$

In the previous expressions, the linearized barycentric velocity is

$$\mathbf{v}_b = \frac{\rho_0 \mathbf{v} + \rho'_0 \mathbf{v}'}{\rho_0 + \rho'_0} = C_0 \mathbf{v} + C'_0 \mathbf{v}'. \quad (\text{B5})$$

Furthermore, $\delta \mathbf{a}$ is the linearized body-force acceleration, which is the deviation of \mathbf{a} from its equilibrium value \mathbf{a}_0 . Expression (19) gives

$$\mathbf{a}_0 = -c_T^2 (\alpha_0 \nabla \rho'_0 + \alpha_1 \Delta \nabla \rho'_0), \quad \delta \mathbf{a} = -c_T^2 (\alpha_0 \nabla \delta \rho' + \alpha_1 \Delta \nabla \delta \rho'). \quad (\text{B6})$$

Equation (B1) can now be written in terms of the deviation of the moments as

$$\partial_t \delta m_a + \sum_b \partial_{ab} \delta m_b = \lambda_a (\delta_{a0} + \delta_{aj} - 1) \delta m_a + \delta \Delta_a + \delta \Phi_a + \xi_a, \quad (\text{B7})$$

with

$$\delta \Delta_a = \frac{\rho_0 \rho'_0}{\rho_{t0}} \lambda_d \delta_a \cdot (\mathbf{v}' - \mathbf{v}) = \lambda_d \delta_a \cdot (C_0 \mathbf{j}' - C'_0 \mathbf{j}), \quad (\text{B8})$$

$$\delta \Phi_a = -c_T^2 \rho_0 \delta_a \cdot (\alpha_0 \nabla \delta \rho' + \alpha_1 \Delta \nabla \delta \rho') + \delta_a \cdot \mathbf{a}_0 \delta \rho + \tau_a : \mathbf{a}_0 \mathbf{j} + \dots \quad (\text{B9})$$

In the above equations, $\mathbf{j} = \rho_0 \mathbf{v}$ is the linearized momentum density, while

$$\begin{aligned} \delta_{aj} &= \sum_{\alpha=1}^D \delta_{a\alpha}, \\ (\delta_a)_\alpha &= \delta_{a\alpha}, \\ (\tau_a)_{\alpha\beta} &= (1 + \delta_{\alpha\beta}) \delta_{a\theta_{\alpha\beta}}. \end{aligned} \quad (\text{B10})$$

Notice that for any vector \mathbf{V} we can write $\delta_a \cdot \mathbf{V} = \delta_{aj} V_a$. Also, $\delta_a \cdot \delta_b = \delta_{aj} \delta_{ab}$, while $\delta \rho = \sum_a \delta_{a0} \delta m_a$ and $\mathbf{j} = \rho_0 \mathbf{v} = \sum_a \delta_a \delta m_a$. We next introduce the Fourier transform as

$$\hat{f}(\mathbf{k}) = \frac{1}{(2\pi)^{D/2}} \int d\mathbf{r} f(\mathbf{r}) e^{-i\mathbf{k}\cdot\mathbf{r}}, \quad f(\mathbf{r}) = \frac{1}{(2\pi)^{D/2}} \int d\mathbf{k} \hat{f}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}}. \quad (\text{B11})$$

According to (40), the fluctuating deviation in Fourier space of the a th moment from its averaged asymptotic value is $\delta \hat{m}_a(\mathbf{k}, t) = \hat{m}_a(\mathbf{k}, t) - \delta_{a0} \hat{\rho}_0(\mathbf{k})$. By Fourier transforming Eq. (B7) we obtain

$$\partial_t \delta \hat{m}_a + \sum_b A_{ab} \delta \hat{m}_b = \lambda_a (\delta_{a0} + \delta_{aj} - 1) \delta \hat{m}_a + \delta \hat{\Delta}_a + \delta \hat{\Phi}_a + \hat{\xi}_a, \quad (\text{B12})$$

where

$$A_{ab}(\mathbf{k}) = \frac{i}{N_b} \int d\mathbf{c} \omega(\mathbf{c}) T_a(\mathbf{c}) T_b(\mathbf{c}) \mathbf{c} \cdot \mathbf{k} \quad (\text{B13})$$

is the advection operator. Furthermore,

$$\delta \hat{\Delta}_a = \lambda_d \delta_a \cdot (\hat{C}_0 * \hat{\mathbf{j}}' - \hat{C}'_0 * \hat{\mathbf{j}}) = - \sum_b \lambda_d \delta_{aj} \delta_{ab} (\hat{C}'_0 * \delta \hat{m}_b - \hat{C}_0 * \delta \hat{m}'_b), \quad (\text{B14})$$

$$\begin{aligned} \delta \hat{\Phi}_a &= -\delta_a \cdot \hat{\rho}_0 * \alpha \delta \hat{\rho}' + \delta_a \cdot \hat{\mathbf{a}}_0 * \delta \hat{\rho} + \tau_a : \hat{\mathbf{a}}_0 * \hat{\mathbf{j}} + \dots, \\ &= - \sum_b (\delta_a \cdot \delta_{b0} \hat{\rho}_0 * \alpha \delta \hat{m}'_b - \delta_a \cdot \delta_{b0} \hat{\mathbf{a}}_0 * \delta \hat{m}_b - \tau_a : \delta_b \hat{\mathbf{a}}_0 * \delta \hat{m}_b + \dots), \end{aligned} \quad (\text{B15})$$

where the asterisk * denotes a normalized convolution in the Fourier space, which is

$$(\hat{g} * \hat{f})(\mathbf{k}) = \frac{1}{(2\pi)^{D/2}} \int d\mathbf{q} \hat{g}(\mathbf{k} - \mathbf{q}) \hat{f}(\mathbf{q}), \quad (\text{B16})$$

while

$$(\alpha \hat{f})(\mathbf{k}) = i c_T^2 \mathbf{k} \alpha(\mathbf{k}) \hat{f}(\mathbf{k}), \quad (\text{B17})$$

$\alpha(\mathbf{k})$ being defined as in Eq. (58). By introducing the indices of species, Eq. (B12) can be written as Eq. (42), with the integral kernel of the time-evolution matrix operator given by

$$\begin{aligned} \mathcal{L}_{ab}^{\sigma\sigma'}(\mathbf{k}, \mathbf{k}') &= \lambda_a (1 - \delta_{a0} - \delta_{aj}) \delta_{ab} \delta(\mathbf{k} - \mathbf{k}') + (2\pi)^{-D/2} \lambda_d \delta_{aj} \delta_{ab} \hat{C}'_0(\mathbf{k} - \mathbf{k}') + A_{ab}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}') \\ &\quad - (2\pi)^{-D/2} [\delta_a \cdot \delta_{b0} \hat{\mathbf{a}}_0(\mathbf{k} - \mathbf{k}') + \tau_a : \delta_b \hat{\mathbf{a}}_0(\mathbf{k} - \mathbf{k}') + \dots], \\ \mathcal{L}_{ab}^{\sigma\sigma'}(\mathbf{k}, \mathbf{k}') &= -(2\pi)^{-D/2} \lambda_d \delta_{aj} \delta_{ab} \hat{C}_0(\mathbf{k} - \mathbf{k}') + (2\pi)^{-D/2} i c_T^2 \delta_a \cdot \delta_{b0} \mathbf{k}' \hat{\rho}_0(\mathbf{k} - \mathbf{k}') \alpha(\mathbf{k}'). \end{aligned} \quad (\text{B18})$$

In particular, in the case of homogeneous equilibrium, $\rho_0(\mathbf{r}) = \rho_0 = \text{const.}$ and from (B6) we find $\mathbf{a}_0 = \mathbf{0}$. As a consequence, $\hat{\rho}_0(\mathbf{k}) = (2\pi)^{D/2} \rho_0 \delta(\mathbf{k})$, $\hat{\mathbf{a}}_0(\mathbf{k}) = \mathbf{0}$. Furthermore, $\hat{C}_0(\mathbf{k}) = (2\pi)^{D/2} C_0 \delta(\mathbf{k})$ and the expressions (B18) become of the form $\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = L_{ab}^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}')$, with

$$\begin{aligned} L_{ab}^{\sigma\sigma}(\mathbf{k}) &= \lambda_a(1 - \delta_{a0} - \delta_{aj})\delta_{ab} + \lambda_d \delta_{aj} \delta_{ab} C'_0 + A_{ab}(\mathbf{k}), \\ L_{ab}^{\sigma\zeta}(\mathbf{k}) &= -\lambda_d \delta_{aj} \delta_{ab} C_0 + i c_\tau^2 \delta_a \cdot \delta_{b0} \mathbf{k} \rho_0 \alpha(\mathbf{k}). \end{aligned} \quad (\text{B19})$$

APPENDIX C: EQUILIBRIUM CORRELATIONS

In this Appendix, we provide expressions for the equilibrium correlations $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ of the lattice Boltzmann modes and their relation to the density and momentum structure factors. It is convenient here to consider the general case of a multicomponent system and maintain the species indices σ, ζ , etc. The structure factors are then defined by

$$\langle \delta \hat{\rho}^\sigma(\mathbf{k}) \delta \hat{\rho}^\zeta(-\mathbf{k}') \rangle = \mathcal{S}_{\rho^\sigma, \rho^\zeta}(\mathbf{k}, \mathbf{k}'), \quad \langle \delta \hat{\mathbf{j}}^\sigma(\mathbf{k}) \delta \hat{\mathbf{j}}^\zeta(-\mathbf{k}') \rangle = \mathcal{S}_{\mathbf{j}^\sigma, \mathbf{j}^\zeta}(\mathbf{k}, \mathbf{k}'), \quad \langle \delta \hat{\mathbf{j}}^\sigma(\mathbf{k}) \delta \hat{\rho}^\zeta(-\mathbf{k}') \rangle = \mathcal{S}_{\mathbf{j}^\sigma, \rho^\zeta}(\mathbf{k}, \mathbf{k}'). \quad (\text{C1})$$

In particular, $\mathcal{S}_{\rho^\sigma, \rho^\zeta}(\mathbf{k}, \mathbf{k}') = \langle |\delta \hat{\rho}^\sigma(\mathbf{k})|^2 \rangle$ is the density structure factor for the σ th species. In the following, we first provide an expression for the correlations $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ defined in Eq. (45). Next, the relations between the structure functions and the correlation matrix are easily obtained as

$$\mathcal{S}_{\rho^\sigma, \rho^\zeta} = \mathcal{G}_{00}^{\sigma\zeta}, \quad \mathcal{S}_{j_a^\sigma, j_b^\zeta} = \mathcal{G}_{\alpha\beta}^{\sigma\zeta}, \quad \mathcal{S}_{j_a^\sigma, \rho^\zeta} = \mathcal{G}_{\alpha 0}^{\sigma\zeta}, \quad (\text{C2})$$

since $\delta \hat{\rho}^\sigma(\mathbf{k}) = \delta \hat{m}_0^\sigma(\mathbf{k})$ and $\delta \hat{j}_a^\sigma(\mathbf{k}) = \delta \hat{m}_a^\sigma(\mathbf{k})$. In order to give an expression for $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$, we write [12,13]

$$\begin{aligned} \langle \delta f(\mathbf{c}, \mathbf{r}) \delta f(\mathbf{c}', \mathbf{r}') \rangle &= \langle [f(\mathbf{c}, \mathbf{r}) - f^{\text{eq}}(\mathbf{c}, \mathbf{r})][f(\mathbf{c}', \mathbf{r}') - f^{\text{eq}}(\mathbf{c}', \mathbf{r}')] \rangle \\ &= \langle f(\mathbf{c}, \mathbf{r}) f(\mathbf{c}', \mathbf{r}') \rangle - f^{\text{eq}}(\mathbf{c}, \mathbf{r}) f^{\text{eq}}(\mathbf{c}', \mathbf{r}') \\ &= \mu f^{\text{eq}}(\mathbf{c}, \mathbf{r}) \delta(\mathbf{c} - \mathbf{c}') \delta(\mathbf{r} - \mathbf{r}') + f_2^{\text{eq}}(\mathbf{c}, \mathbf{r}, \mathbf{c}', \mathbf{r}') - f^{\text{eq}}(\mathbf{c}, \mathbf{r}) f^{\text{eq}}(\mathbf{c}', \mathbf{r}'), \end{aligned} \quad (\text{C3})$$

where $f_2^{\text{eq}}(\mathbf{c}, \mathbf{r}, \mathbf{c}', \mathbf{r}') = f_2^{\text{eq}}(\mathbf{c}', \mathbf{r}', \mathbf{c}, \mathbf{r})$ is the averaged two-point distribution function. Here $f(\mathbf{c}, \mathbf{r})$ is the value reached by the Boltzmann distribution function $f(\mathbf{c}, \mathbf{r}, t)$ asymptotically for $t \rightarrow \infty$, while $\delta f(\mathbf{c}, \mathbf{r})$ is the fluctuation of $f(\mathbf{c}, \mathbf{r})$ from the equilibrium Maxwellian (Fig. 1), $f^{\text{eq}}(\mathbf{c}, \mathbf{r}) = f^{\text{eq}}(\rho_0(\mathbf{r}), \mathbf{0}; \mathbf{c})$, with $\rho_0(\mathbf{r})$ the averaged mass density at equilibrium. In the first line in Eq. (C3), only the definition of the fluctuating deviation from the equilibrium distribution is used. The last line, on the other hand, can be obtained by assuming a Gaussian character of the fluctuations of the Boltzmann distribution function and requiring that its first and second moments are identical to the corresponding moments of the exact N -particle phase-space density, as introduced by Klimontovich [33]. In this way, two-particle correlations mediated by nonideal fluid forces are maintained, while possible higher-order, non-Gaussian correlations are neglected. The first term in the last line of Eq. (C3) is the self part of $\langle f(\mathbf{c}, \mathbf{r}) f(\mathbf{c}', \mathbf{r}') \rangle$, whereas $f_2^{\text{eq}}(\mathbf{c}, \mathbf{r}, \mathbf{c}', \mathbf{r}')$ is its distinct part, also known as the reduced two-particle phase-space density distribution [42]. For a multicomponent fluid at equilibrium, a natural generalization of Eq. (C3) can be proposed as

$$\langle \delta f^\sigma(\mathbf{c}, \mathbf{r}) \delta f^\zeta(\mathbf{c}', \mathbf{r}') \rangle = \mu^\sigma f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) \delta(\mathbf{c} - \mathbf{c}') \delta(\mathbf{r} - \mathbf{r}') \delta_{\sigma\zeta} + f_2^{\text{eq}, \sigma\zeta}(\mathbf{c}, \mathbf{r}, \mathbf{c}', \mathbf{r}') - f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) f^{\text{eq}, \zeta}(\mathbf{c}', \mathbf{r}'), \quad (\text{C4})$$

where $f_2^{\text{eq}, \sigma\zeta}(\mathbf{c}, \mathbf{r}, \mathbf{c}', \mathbf{r}') = f_2^{\text{eq}, \zeta\sigma}(\mathbf{c}', \mathbf{r}', \mathbf{c}, \mathbf{r})$ is the averaged two-point multicomponent distribution function, while $f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) = f^{\text{eq}}(\rho_0^\sigma(\mathbf{r}), \mathbf{0}; \mathbf{c})$, with $\rho_0^\sigma(\mathbf{r})$ the averaged σ th mass density at equilibrium. Furthermore, we introduce the *pair correlation function* $\gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}')$ using the ansatz [42]

$$f_2^{\text{eq}, \sigma\zeta}(\mathbf{c}, \mathbf{r}, \mathbf{c}', \mathbf{r}') = f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) f^{\text{eq}, \zeta}(\mathbf{c}', \mathbf{r}') [1 + \gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}')], \quad (\text{C5})$$

where $\gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') = \gamma^{\zeta\sigma}(\mathbf{r}', \mathbf{r})$. Note that this definition of the pair correlation function differs from the usual definitions of the pair distribution function $g^{\sigma\zeta}(\mathbf{r}, \mathbf{r}')$ by unity, i.e., $\gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') = g^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') - 1$ [42]. As a consequence, $\gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') \rightarrow 0$ in the limit of $|\mathbf{r} - \mathbf{r}'| \rightarrow \infty$. Using the thus introduced pair correlation function, Eq. (C4) can be written as

$$\langle \delta f^\sigma(\mathbf{c}, \mathbf{r}) \delta f^\zeta(\mathbf{c}', \mathbf{r}') \rangle = \mu^\sigma f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) \delta(\mathbf{c} - \mathbf{c}') \delta(\mathbf{r} - \mathbf{r}') \delta_{\sigma\zeta} + f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) f^{\text{eq}, \zeta}(\mathbf{c}', \mathbf{r}') \gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}'). \quad (\text{C6})$$

The presence of the pair correlation function is directly related to the self-generated body force. The choice $\gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') = 0$ indeed corresponds to an ideal mixture. In our case,

$$\langle \delta f^\sigma(\mathbf{c}, \mathbf{r}) \delta f^\zeta(\mathbf{c}', \mathbf{r}') \rangle = \mu \omega(\mathbf{c}) \rho_0^\sigma(\mathbf{r}) \delta(\mathbf{c} - \mathbf{c}') \delta(\mathbf{r} - \mathbf{r}') \delta_{\sigma\zeta} + \omega(\mathbf{c}) \omega(\mathbf{c}') \rho_0^\sigma(\mathbf{r}) \rho_0^\zeta(\mathbf{r}') \gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}'), \quad (\text{C7})$$

where the expression $f^{\text{eq}, \sigma}(\mathbf{c}, \mathbf{r}) = f^{\text{eq}}(\rho_0^\sigma(\mathbf{r}), \mathbf{0}; \mathbf{c}) = \omega(\mathbf{c}) \rho_0^\sigma(\mathbf{r})$ for the equilibrium Maxwellian has been used, while $\mu^\sigma = \mu$ for each species. Equation (C7) is an important input to a Boltzmann-Langevin model and specifies its complete structure of the equilibrium correlations. In particular, Eq. (C7) encapsulates also the equilibrium correlations of the nonhydrodynamic modes, which are coupled to hydrodynamic modes at finite length scales. In our case, the statistics of the nonhydrodynamic modes is

the same as for an ideal gas [12,13,35]. By expressing the previous equation in terms of the moments we obtain

$$\langle \delta m_a^\sigma(\mathbf{r}) \delta m_b^\zeta(\mathbf{r}') \rangle = \mu \rho_0^\sigma(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}') N_a \delta_{ab} \delta_{\sigma\zeta} + \rho_0^\sigma(\mathbf{r}) \rho_0^\zeta(\mathbf{r}') \gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') \delta_{a0} \delta_{b0}, \quad (\text{C8})$$

which, after Fourier-transforming, becomes [see Eq. (45)]

$$\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = \langle \delta \hat{m}_a^\sigma(\mathbf{k}) \delta \hat{m}_b^\zeta(-\mathbf{k}') \rangle = (2\pi)^{-D/2} \mu \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') N_a \delta_{ab} \delta_{\sigma\zeta} + (\hat{\rho}_0^\sigma \hat{\rho}_0^\zeta * \hat{\gamma}^{\sigma\zeta})(\mathbf{k}, -\mathbf{k}') \delta_{a0} \delta_{b0}, \quad (\text{C9})$$

where the double asterisk $*$ denotes a normalized biconvolution in the Fourier space, which is

$$(\hat{g} * \hat{f})(\mathbf{k}, \mathbf{k}') = \frac{1}{(2\pi)^D} \int d\mathbf{q} d\mathbf{q}' \hat{g}(\mathbf{k} - \mathbf{q}, \mathbf{k}' - \mathbf{q}') \hat{f}(\mathbf{q}, \mathbf{q}'), \quad (\text{C10})$$

while

$$\hat{\gamma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = \frac{1}{(2\pi)^D} \int d\mathbf{r} d\mathbf{r}' \gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') e^{i\mathbf{k}\cdot\mathbf{r} + i\mathbf{k}'\cdot\mathbf{r}'}. \quad (\text{C11})$$

Notice that $\hat{\gamma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = \hat{\gamma}^{\zeta\sigma}(\mathbf{k}', \mathbf{k})$. Based on Eq. (C2), we finally obtain the expression of the relevant structure factors in terms of the Fourier transform of the pair correlation function. For the mass density they read

$$\mathcal{S}_{\rho^\sigma, \rho^\zeta}(\mathbf{k}, \mathbf{k}') = (2\pi)^{-D/2} \frac{k_B T}{c_T^2} \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') \delta_{\sigma\zeta} + (\hat{\rho}_0^\sigma \hat{\rho}_0^\zeta * \hat{\gamma}^{\sigma\zeta})(\mathbf{k}, -\mathbf{k}'), \quad (\text{C12})$$

where we used $\mu = k_B T / c_T^2$, while for the momentum they are

$$\mathcal{S}_{j^\sigma, j^\zeta}(\mathbf{k}, \mathbf{k}') = (2\pi)^{-D/2} k_B T \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') \delta_{\sigma\zeta} \mathbb{1}, \quad \mathcal{S}_{j^\sigma, \rho^\zeta}(\mathbf{k}, \mathbf{k}') = \mathbf{0}, \quad (\text{C13})$$

$\mathbb{1}$ being the $D \times D$ identity. Specializing to the homogeneous case, $\rho^\sigma(\mathbf{r}) = \rho^\sigma = \text{const.}$ and hence $\hat{\rho}_0^\sigma(\mathbf{k}) = (2\pi)^{D/2} \delta(\mathbf{k})$. Furthermore, translational invariance of the left-hand side of Eq. (C5) implies $\gamma^{\sigma\zeta}(\mathbf{r}, \mathbf{r}') = \gamma^{\sigma\zeta}(\mathbf{r} - \mathbf{r}')$ on the right-hand side. Thus, we have $\hat{\gamma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = (2\pi)^{D/2} \hat{\gamma}^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}')$ in Eq. (C11). Notice that $\hat{\gamma}^{\sigma\zeta}(\mathbf{k}) = \hat{\gamma}^{\zeta\sigma}(-\mathbf{k})$. Equation (C9) then becomes $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = G_{ab}^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}')$, with

$$G_{ab}^{\sigma\zeta}(\mathbf{k}) = \mu \rho^\sigma N_a \delta_{ab} \delta_{\sigma\zeta} + (2\pi)^{D/2} \rho^\sigma \rho^\zeta \hat{\gamma}^{\sigma\zeta}(\mathbf{k}) \delta_{a0} \delta_{b0}. \quad (\text{C14})$$

We can thus extract the relevant information from the diagonal part in Fourier space. By writing the generic structure factor as $\mathcal{S}(\mathbf{k}, \mathbf{k}') = S(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}')$, the equivalent of Eqs. (C12) and (C13) are

$$S_{\rho^\sigma, \rho^\zeta}(\mathbf{k}) = \frac{k_B T}{c_T^2} \rho^\sigma \delta_{\sigma\zeta} + (2\pi)^{D/2} \rho^\sigma \rho^\zeta \hat{\gamma}^{\sigma\zeta}(\mathbf{k}) \quad (\text{C15})$$

and

$$\mathcal{S}_{j^\sigma, j^\zeta}(\mathbf{k}) = k_B T \rho^\sigma \delta_{\sigma\zeta} \mathbb{1}, \quad \mathcal{S}_{j^\sigma, \rho^\zeta}(\mathbf{k}) = \mathbf{0}. \quad (\text{C16})$$

APPENDIX D: CALCULATION OF NOISE COVARIANCES

In this Appendix we detail the calculation for the noise covariances in both the homogeneous and the nonhomogeneous cases. Also here we consider the general case of a multicomponent system with species indices σ, ζ, κ , etc. Furthermore, we set

$$r_a^\sigma = \lambda_a^\sigma (1 - \delta_{a0} - \delta_{aj}), \quad d_a^{\zeta\sigma} = \lambda_d \delta_{aj} (\delta_{\sigma\zeta} - C_0^\sigma), \quad \alpha^{\sigma\zeta} = (1 - \delta_{\sigma\zeta}) \alpha, \quad (\text{D1})$$

for short. The following identities will be used:

$$N_a A_{ba} = N_b A_{ab}, \quad A_{a0}(\mathbf{k}) = i c_T^2 \boldsymbol{\delta}_a \cdot \mathbf{k}, \quad d_a^{\zeta\sigma} = \delta_{aj} d_a^{\zeta\sigma}, \quad \rho_0^\zeta d_a^{\zeta\sigma} = \rho_0^\sigma d_a^{\zeta\sigma}, \quad \alpha^{\sigma\zeta}(\mathbf{k}) = \alpha^{\zeta\sigma}(-\mathbf{k}). \quad (\text{D2})$$

The expressions of $d_a^{\zeta\sigma}$ and $\alpha^{\sigma\zeta}$ given here depend on the model used. Nevertheless, the results of the present section are valid for any multicomponent model whose time-evolution operator has one of the forms given in Eqs. (D3) or (D13), provided the corresponding $d_a^{\zeta\sigma}$ and $\alpha^{\sigma\zeta}$ satisfy the identities given above. We remark that the species are assumed to be non-self-interacting. The inclusion of the self-interaction is straightforward and does not lead to any change in the noise covariance.

1. Homogeneous equilibrium

The time-evolution matrix $L_{ab}^{\sigma\zeta}(\mathbf{k})$ in (B19) can be written as

$$L_{ab}^{\sigma\zeta}(\mathbf{k}) = r_a^\sigma \delta_{ab} \delta_{\sigma\zeta} + d_a^{\zeta\sigma} \delta_{ab} + \delta_{\sigma\zeta} A_{ab}(\mathbf{k}) + i c_T^2 \boldsymbol{\delta}_a \cdot \delta_{b0} \mathbf{k} \rho_0^\sigma \alpha^{\sigma\zeta}(\mathbf{k}). \quad (\text{D3})$$

Based on this expression, we need to use the result for the noise covariances $\Xi_{ab}^{\sigma\zeta}(\mathbf{k})$ given by Eq. (51). Using $G_{ab}^{\sigma\zeta}(\mathbf{k})$ given by Eq. (C14), we obtain

$$\begin{aligned}\Xi_{ab}^{\sigma\zeta}(\mathbf{k}) &= \sum_{c,\kappa} G_{ac}^{\sigma\kappa}(\mathbf{k})L_{bc}^{\zeta\kappa}(-\mathbf{k}) + \sum_{c,\kappa} L_{ac}^{\sigma\kappa}(\mathbf{k})G_{cb}^{\zeta\kappa}(\mathbf{k}) \\ &= \mu[\rho_0^\sigma N_a L_{ba}^{\zeta\sigma}(-\mathbf{k}) + L_{ab}^{\zeta\sigma}(\mathbf{k})\rho_0^\zeta N_b] + (2\pi)^{D/2} \sum_{\kappa} [\rho_0^\sigma \rho_0^\kappa \hat{\gamma}^{\sigma\kappa}(\mathbf{k})\delta_{a0}L_{b0}^{\zeta\kappa}(-\mathbf{k}) + L_{a0}^{\sigma\kappa}(\mathbf{k})\rho_0^\kappa \rho_0^\zeta \hat{\gamma}^{\kappa\zeta}(\mathbf{k})\delta_{b0}] \\ &= 2\mu\rho_0^\sigma N_a (r_a^\sigma \delta_{\sigma\zeta} + c_\tau^2 d_a^{\zeta\sigma})\delta_{ab} - i\rho_0^\sigma \rho_0^\zeta (\delta_{a0}\delta_b - \delta_{b0}\delta_a) \cdot \mathbf{k} [\mu\alpha^{\sigma\zeta}(\mathbf{k}) + (2\pi)^{D/2} \hat{\gamma}^{\sigma\zeta}(\mathbf{k})] \\ &\quad - (2\pi)^{D/2} i c_\tau \rho_0^\sigma \rho_0^\zeta \left[\delta_{a0}\delta_b \cdot \mathbf{k} \sum_{\kappa} \rho_0^\kappa \hat{\gamma}^{\sigma\kappa}(\mathbf{k})\alpha^{\zeta\kappa}(\mathbf{k}) - \delta_{b0}\delta_a \cdot \mathbf{k} \sum_{\kappa} \rho_0^\kappa \alpha^{\sigma\kappa}(\mathbf{k})\hat{\gamma}^{\kappa\zeta}(\mathbf{k}) \right],\end{aligned}\quad (\text{D4})$$

which can be written as

$$\Xi_{ab}^{\sigma\zeta}(\mathbf{k}) = 2\mu\rho_0^\sigma N_a (r_a^\sigma \delta_{\sigma\zeta} + d_a^{\zeta\sigma})\delta_{ab} + \delta_{a0}\delta_b \cdot \boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}) + \delta_{b0}\delta_a \cdot \boldsymbol{\Sigma}^{\zeta\sigma}(-\mathbf{k}), \quad (\text{D5})$$

with

$$\boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}) = -i c_\tau^2 \rho_0^\sigma \rho_0^\zeta \mathbf{k} \left[\mu\alpha^{\sigma\zeta}(\mathbf{k}) + (2\pi)^{D/2} \sum_{\kappa} \rho_0^\kappa \hat{\gamma}^{\sigma\kappa}(\mathbf{k})\alpha^{\zeta\kappa}(\mathbf{k}) + (2\pi)^{D/2} \hat{\gamma}^{\sigma\zeta}(\mathbf{k}) \right]. \quad (\text{D6})$$

Noting that $\Xi_{00}^{\sigma\zeta}(\mathbf{k}) = 0$, we necessarily have to set $\xi_0^\sigma(\mathbf{k}, t) = 0$ identically. It follows that all correlations of the form $\langle \xi_0^\sigma(\mathbf{k}, t)\xi_b^\zeta(-\mathbf{k}', t') \rangle$ [or equivalently $\langle \xi_a^\sigma(\mathbf{k}, t)\xi_0^\zeta(-\mathbf{k}', t') \rangle$] must vanish. To be self-consistent, we then impose $\Xi_{0b}^{\sigma\zeta}(\mathbf{k}) = 0$ for any b [or equivalently $\Xi_{a0}^{\sigma\zeta}(\mathbf{k}) = 0$ for any a], obtaining $\boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}) = \mathbf{0}$ [or equivalently $\boldsymbol{\Sigma}^{\zeta\sigma}(-\mathbf{k}) = \mathbf{0}$]. As a consequence, from (D5) we obtain

$$\Xi_{ab}^{\sigma\zeta}(\mathbf{k}) = 2\mu\rho_0^\sigma N_a (r_a^\sigma \delta_{\sigma\zeta} + d_a^{\zeta\sigma})\delta_{ab}, \quad (\text{D7})$$

which is independent of \mathbf{k} . By Fourier-transforming back to real space and using $\mu = k_B T / c_\tau^2$, Eq. (D7) yields the following noise correlations:

$$\langle \xi_a^\sigma(\mathbf{r}, t)\xi_b^\zeta(\mathbf{r}', t') \rangle = 2 \frac{k_B T}{c_\tau^2} \rho_0 N_a (r_a^\sigma \delta_{\sigma\zeta} + d_a^{\zeta\sigma})\delta_{ab}\delta(\mathbf{r} - \mathbf{r}')\delta(t - t'). \quad (\text{D8})$$

In particular, in our case we have

$$\langle \xi_a^\sigma(\mathbf{r}, t)\xi_b^\zeta(\mathbf{r}', t') \rangle = 2 \frac{k_B T}{c_\tau^2} \rho_0 N_a \lambda_a [(1 - \delta_{a0})\delta_{\sigma\zeta} - C_0^\zeta \delta_{aj}]\delta_{ab}\delta(\mathbf{r} - \mathbf{r}')\delta(t - t'), \quad (\text{D9})$$

which results in Eqs. (53). Furthermore, the self-consistency condition $\boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}) = \mathbf{0}$ for any \mathbf{k} leads to

$$\mu\alpha^{\sigma\zeta} + (2\pi)^{D/2} \sum_{\kappa} \rho_0^\kappa \hat{\gamma}^{\sigma\kappa} \alpha^{\zeta\kappa} + (2\pi)^{D/2} \hat{\gamma}^{\sigma\zeta} = 0. \quad (\text{D10})$$

In particular, in our case we have

$$\begin{aligned}\rho_0' \hat{\Gamma} \alpha + \hat{\gamma} &= 0, \\ \mu\alpha + (2\pi)^{D/2} \rho_0 \hat{\gamma} \alpha + (2\pi)^{D/2} \hat{\Gamma} &= 0.\end{aligned}\quad (\text{D11})$$

Such a system can be easily solved as

$$\begin{aligned}(2\pi)^{D/2} \hat{\gamma} &= \frac{\mu\rho_0' \alpha^2}{1 - \rho_0 \rho_0' \alpha^2}, \\ (2\pi)^{D/2} \hat{\Gamma} &= -\frac{\mu\alpha}{1 - \rho_0 \rho_0' \alpha^2},\end{aligned}\quad (\text{D12})$$

which inserted in Eqs. (54) lead to Eqs. (57).

2. Nonhomogeneous equilibrium

The integral kernel $\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ in (B18) can be written as

$$\begin{aligned}\mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') &= r_a^\sigma \delta_{ab} \delta_{\sigma\zeta} \delta(\mathbf{k} - \mathbf{k}') + (2\pi)^{-D/2} \delta_{ab} \hat{d}_a^{\sigma\zeta}(\mathbf{k} - \mathbf{k}') + \delta_{\sigma\zeta} A_{ab}(\mathbf{k})\delta(\mathbf{k} - \mathbf{k}') \\ &\quad - (2\pi)^{-D/2} \delta_{\sigma\zeta} [\delta_a \cdot \delta_{b0} \hat{\mathbf{a}}_0^\sigma(\mathbf{k} - \mathbf{k}') + \boldsymbol{\tau}_a : \delta_b \hat{\mathbf{a}}_0^\sigma(\mathbf{k} - \mathbf{k}') + \dots] + (2\pi)^{-D/2} i c_\tau^2 \delta_a \cdot \delta_{b0} \mathbf{k}' \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') \alpha^{\sigma\zeta}(\mathbf{k}').\end{aligned}\quad (\text{D13})$$

Based on this expression, we need to use the general result for the noise covariances $\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ given in Eqs. (44). Using $\mathcal{G}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}')$ given by Eq. (C9), we obtain

$$\begin{aligned}
\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') &= \int d\mathbf{q} \sum_{c,\kappa} \mathcal{G}_{ac}^{\sigma\kappa}(\mathbf{k}, \mathbf{q}) \mathcal{L}_{bc}^{\zeta\kappa}(-\mathbf{k}', -\mathbf{q}) + \int d\mathbf{q} \sum_{c,\kappa} \mathcal{L}_{ac}^{\sigma\kappa}(\mathbf{k}, \mathbf{q}) \mathcal{G}_{cb}^{\zeta\kappa}(\mathbf{q}, \mathbf{k}') \\
&= (2\pi)^{-D/2} \mu \int d\mathbf{q} [\rho_0^\sigma(\mathbf{k} - \mathbf{q}) N_a \mathcal{L}_{ba}^{\zeta\sigma}(-\mathbf{k}', -\mathbf{q}) + \mathcal{L}_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{q}) \rho_0^\zeta(\mathbf{q} - \mathbf{k}') N_b] \\
&\quad + \int d\mathbf{q} \sum_{\kappa} [(\hat{\rho}_0^\sigma \hat{\rho}_0^\kappa * \hat{\gamma}^{\sigma\kappa})(\mathbf{k}, -\mathbf{q}) \delta_{a0} \mathcal{L}_{b0}^{\zeta\kappa}(-\mathbf{k}', -\mathbf{q}) + \mathcal{L}_{a0}^{\sigma\kappa}(\mathbf{k}, \mathbf{q}) (\hat{\rho}_0^\kappa \hat{\rho}_0^\zeta * \hat{\gamma}^{\kappa\zeta})(\mathbf{q}, -\mathbf{k}') \delta_{b0}] \\
&= (2\pi)^{-D/2} 2\mu N_a [\hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') r_a^\sigma \delta_{\sigma\zeta} + (\hat{\rho}_0^\sigma * \hat{d}_a^{\zeta\sigma})(\mathbf{k} - \mathbf{k}')] \delta_{ab} + (2\pi)^{-D/2} \mu \delta_{\sigma\zeta} N_b A_{ab}(\mathbf{k} - \mathbf{k}') \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') \\
&\quad - (2\pi)^{-D/2} \mu \delta_{\sigma\zeta} [\delta_b \cdot \delta_{a0} (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma)(\mathbf{k} - \mathbf{k}') + c_T^2 \boldsymbol{\tau}_b : \delta_a (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma)(\mathbf{k} - \mathbf{k}') + \dots] \\
&\quad - (2\pi)^{-D/2} \mu \delta_{\sigma\zeta} [\delta_a \cdot \delta_{b0} (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma)(\mathbf{k} - \mathbf{k}') + c_T^2 \boldsymbol{\tau}_a : \delta_b (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma)(\mathbf{k} - \mathbf{k}') + \dots] \\
&\quad - (2\pi)^{-D} i c_T^2 \mu (\delta_{a0} \delta_b - \delta_{b0} \delta_a) \cdot \int d\mathbf{q} \mathbf{q} \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}') \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}) \alpha^{\sigma\zeta}(\mathbf{q}) \\
&\quad - i c_T^2 (\delta_{a0} \delta_b \cdot \mathbf{k}' - \delta_{b0} \delta_a \cdot \mathbf{k}) (\hat{\rho}_0^\sigma \hat{\rho}_0^\zeta * \hat{\gamma}^{\sigma\zeta})(\mathbf{k}, -\mathbf{k}') \\
&\quad - \delta_{a0} \delta_b \cdot [\hat{\rho}_0^\sigma (\hat{\rho}_0^\zeta * \hat{\mathbf{a}}_0^\zeta) * \hat{\gamma}^{\sigma\zeta}](\mathbf{k}, -\mathbf{k}') - \delta_{b0} \delta_a \cdot [(\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma) \hat{\rho}_0^\zeta * \hat{\gamma}^{\sigma\zeta}](\mathbf{k}, -\mathbf{k}') \\
&\quad - (2\pi)^{-D/2} i c_T^2 \delta_{a0} \delta_b \cdot \int d\mathbf{q} \mathbf{q} \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}') \sum_{\kappa} \alpha^{\zeta\kappa}(\mathbf{q}) (\hat{\rho}_0^\sigma \hat{\rho}_0^\kappa * \hat{\gamma}^{\sigma\kappa})(\mathbf{k}, -\mathbf{q}) \\
&\quad + (2\pi)^{-D/2} i c_T^2 \delta_{b0} \delta_a \cdot \int d\mathbf{q} \mathbf{q} \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}) \sum_{\kappa} \alpha^{\sigma\kappa}(\mathbf{q}) (\hat{\rho}_0^\kappa \hat{\rho}_0^\zeta * \hat{\gamma}^{\kappa\zeta})(\mathbf{q}, -\mathbf{k}'), \tag{D14}
\end{aligned}$$

which can be written in a more compact form as

$$\begin{aligned}
\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') &= (2\pi)^{-D/2} 2\mu N_a [\hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') r_a^\sigma \delta_{\sigma\zeta} + (\hat{\rho}_0^\sigma * \hat{d}_a^{\zeta\sigma})(\mathbf{k} - \mathbf{k}')] \delta_{ab} + \delta_{a0} \delta_b \cdot \boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') + \delta_{b0} \delta_a \cdot \boldsymbol{\Sigma}^{\zeta\sigma}(-\mathbf{k}', -\mathbf{k}) \\
&\quad + (2\pi)^{-D/2} \mu \delta_{\sigma\zeta} [N_b A_{ab}(\mathbf{k} - \mathbf{k}') \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') - \phi_{ab}^\sigma(\mathbf{k} - \mathbf{k}')], \tag{D15}
\end{aligned}$$

where

$$\phi_{ab}^\sigma = \delta_a \cdot \delta_{b0} (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma) + \delta_b \cdot \delta_{a0} (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma) + c_T^2 [\boldsymbol{\tau}_a : \delta_b (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma) + \boldsymbol{\tau}_b : \delta_a (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma)] + \dots \tag{D16}$$

and

$$\begin{aligned}
\boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') &= -(2\pi)^{-D} i c_T^2 \int d\mathbf{q} \mathbf{q} \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}') \left[\mu \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}) \alpha^{\sigma\zeta}(\mathbf{q}) + (2\pi)^{D/2} \sum_{\kappa} \alpha^{\zeta\kappa}(\mathbf{q}) (\hat{\rho}_0^\sigma \hat{\rho}_0^\kappa * \hat{\gamma}^{\sigma\kappa})(\mathbf{k}, -\mathbf{q}) \right] \\
&\quad - i c_T^2 \mathbf{k}' (\hat{\rho}_0^\sigma \hat{\rho}_0^\zeta * \hat{\gamma}^{\sigma\zeta})(\mathbf{k}, -\mathbf{k}') - [\hat{\rho}_0^\sigma (\hat{\rho}_0^\zeta * \hat{\mathbf{a}}_0^\zeta) * \hat{\gamma}^{\sigma\zeta}](\mathbf{k}, -\mathbf{k}'). \tag{D17}
\end{aligned}$$

Multiplying Eq. (A19) by $\rho_0^\sigma(\mathbf{k})$, we obtain the following relation:

$$N_b A_{ab}(\mathbf{k}) \rho_0^\sigma(\mathbf{k}) = \delta_a \cdot \delta_{b0} i c_T^2 \mathbf{k} \rho_0^\sigma(\mathbf{k}) + \delta_b \cdot \delta_{a0} i c_T^2 \mathbf{k} \rho_0^\sigma(\mathbf{k}) + c_T^2 [\boldsymbol{\tau}_a : \delta_b i c_T^2 \mathbf{k} \rho_0^\sigma(\mathbf{k}) + \boldsymbol{\tau}_b : \delta_a i c_T^2 \mathbf{k} \rho_0^\sigma(\mathbf{k})] + \dots \tag{D18}$$

Furthermore, the equilibrium condition (21) written in Fourier space reads

$$i c_T^2 \mathbf{k} \rho_0^\sigma(\mathbf{k}) = (\hat{\rho}_0^\sigma * \hat{\mathbf{a}}_0^\sigma)(\mathbf{k}). \tag{D19}$$

It follows that $N_b A_{ab} \hat{\rho}_0^\sigma = \phi_{ab}^\sigma$ and Eq. (D15) reduces to

$$\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = (2\pi)^{-D/2} 2\mu N_a [\hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') r_a^\sigma \delta_{\sigma\zeta} + (\hat{\rho}_0^\sigma * \hat{d}_a^{\zeta\sigma})(\mathbf{k} - \mathbf{k}')] \delta_{ab} + \delta_{a0} \delta_b \cdot \boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') + \delta_{b0} \delta_a \cdot \boldsymbol{\Sigma}^{\zeta\sigma}(-\mathbf{k}', -\mathbf{k}). \tag{D20}$$

Again, as in the homogeneous case, we remark that $\Xi_{00}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = 0$ and set $\xi_0^\sigma(\mathbf{k}, t) = 0$. It follows that all the correlations of the form $\langle \xi_0^\sigma(\mathbf{k}, t) \xi_b^\zeta(-\mathbf{k}', t') \rangle$ [or equivalently $\langle \xi_a^\sigma(\mathbf{k}, t) \xi_0^\zeta(-\mathbf{k}', t') \rangle$] must vanish. To be self-consistent, we then impose $\Xi_{0b}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = 0$ for any b [or equivalently $\Xi_{a0}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = 0$ for any a], obtaining $\boldsymbol{\Sigma}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = \mathbf{0}$ [or equivalently $\boldsymbol{\Sigma}^{\zeta\sigma}(-\mathbf{k}', -\mathbf{k}) = \mathbf{0}$]. As a consequence,

$$\Xi_{ab}^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = (2\pi)^{-D/2} 2\mu N_a [\hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{k}') r_a^\sigma \delta_{\sigma\zeta} + (\hat{\rho}_0^\sigma * \hat{d}_a^{\zeta\sigma})(\mathbf{k} - \mathbf{k}')] \delta_{ab}. \tag{D21}$$

By returning to real space and using $\mu = k_B T / c_T^2$, Eq. (D21) gives the following noise correlations:

$$\langle \xi_a^\sigma(\mathbf{r}, t) \xi_b^\zeta(\mathbf{r}', t') \rangle = 2 \frac{k_B T}{c_T^2} \rho_0(\mathbf{r}) N_a [r_a^\sigma \delta_{\sigma\zeta} + d_a^{\zeta\sigma}(\mathbf{r})] \delta_{ab} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t'). \quad (\text{D22})$$

In particular, in our case we have

$$\langle \xi_a^\sigma(\mathbf{r}, t) \xi_b^\zeta(\mathbf{r}', t') \rangle = 2 \frac{k_B T}{c_T^2} \rho_0(\mathbf{r}) N_a \lambda_a [(1 - \delta_{a0}) \delta_{\sigma\zeta} - C_0^\zeta(\mathbf{r}) \delta_{aj}] \delta_{ab} \delta(\mathbf{r} - \mathbf{r}') \delta(t - t'), \quad (\text{D23})$$

which results in Eqs. (60). Furthermore, from Eq. (D19) we can write $(\hat{\rho}_0^\zeta * \hat{\mathbf{a}}_0^\zeta)(\mathbf{q} - \mathbf{k}') = i c_T^2 (\mathbf{q} - \mathbf{k}') \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}')$ and hence,

$$\hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}') (\hat{\rho}_0^\zeta * \hat{\mathbf{a}}_0^\zeta)(\mathbf{q} - \mathbf{k}') \hat{\gamma}^{\sigma\zeta}(\mathbf{q}', -\mathbf{q}) = i c_T^2 (\mathbf{q} - \mathbf{k}') \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}') \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}') \hat{\gamma}^{\sigma\zeta}(\mathbf{q}', -\mathbf{q}). \quad (\text{D24})$$

By integrating over $d\mathbf{q} d\mathbf{q}'$ and rearranging, we obtain

$$i c_T^2 \mathbf{k}' (\hat{\rho}_0^\sigma \hat{\rho}_0^\zeta * \hat{\gamma}^{\sigma\zeta})(\mathbf{k}, -\mathbf{k}') + [\hat{\rho}_0^\sigma (\hat{\rho}_0^\zeta * \hat{\mathbf{a}}_0^\zeta) * \hat{\gamma}^{\sigma\zeta}](\mathbf{k}, -\mathbf{k}') = (2\pi)^{-D} i c_T^2 \int d\mathbf{q} d\mathbf{q}' \mathbf{q} \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}') \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}') \hat{\gamma}^{\sigma\zeta}(\mathbf{q}', -\mathbf{q}) \quad (\text{D25})$$

and hence

$$\begin{aligned} \Sigma^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = & -(2\pi)^{-D} i c_T^2 \int d\mathbf{q} \mathbf{q} \hat{\rho}_0^\zeta(\mathbf{q} - \mathbf{k}') \left[\mu \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}) \alpha^{\sigma\zeta}(\mathbf{q}) + (2\pi)^{D/2} \sum_{\kappa} \alpha^{\zeta\kappa}(\mathbf{q}) (\hat{\rho}_0^\sigma \hat{\rho}_0^\kappa * \hat{\gamma}^{\sigma\kappa})(\mathbf{k}, -\mathbf{q}) \right. \\ & \left. + \int d\mathbf{q}' \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}') \hat{\gamma}^{\sigma\zeta}(\mathbf{q}', -\mathbf{q}) \right]. \end{aligned} \quad (\text{D26})$$

The self-consistency condition $\Sigma^{\sigma\zeta}(\mathbf{k}, \mathbf{k}') = \mathbf{0}$ for any \mathbf{k}' implies the vanishing of the term in the round brackets, leading to

$$\int d\mathbf{q}' \hat{\rho}_0^\sigma(\mathbf{k} - \mathbf{q}') \left[\mu \alpha^{\sigma\zeta}(\mathbf{q}') \delta(\mathbf{q}' - \mathbf{k}') + (2\pi)^{-D/2} \sum_{\kappa} \alpha^{\zeta\kappa}(\mathbf{k}') \int d\mathbf{q} \hat{\rho}_0^\kappa(\mathbf{q} - \mathbf{k}') \hat{\gamma}^{\sigma\kappa}(\mathbf{q}', -\mathbf{q}) + \hat{\gamma}^{\sigma\zeta}(\mathbf{q}', -\mathbf{k}') \right] = 0. \quad (\text{D27})$$

The validity of this condition for any \mathbf{k} again implies the vanishing of the term in the round brackets and hence

$$\mu \alpha^{\sigma\zeta}(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}') + (2\pi)^{-D/2} \sum_{\kappa} \alpha^{\zeta\kappa}(\mathbf{k}') \int d\mathbf{q} \hat{\rho}_0^\kappa(\mathbf{q} - \mathbf{k}') \hat{\gamma}^{\sigma\kappa}(\mathbf{k}, -\mathbf{q}) + \hat{\gamma}^{\sigma\zeta}(\mathbf{k}, -\mathbf{k}') = 0. \quad (\text{D28})$$

In particular, in our case we have

$$\begin{aligned} (2\pi)^{-D/2} \alpha(\mathbf{k}') \int d\mathbf{q} \hat{\rho}'_0(\mathbf{q} - \mathbf{k}') \hat{\Gamma}(\mathbf{k}, -\mathbf{q}) + \hat{\gamma}(\mathbf{k}, -\mathbf{k}') &= 0, \\ \mu \alpha(\mathbf{k}) \delta(\mathbf{k} - \mathbf{k}') + (2\pi)^{-D/2} \alpha(\mathbf{k}') \int d\mathbf{q} \hat{\rho}_0(\mathbf{q} - \mathbf{k}') \hat{\gamma}(\mathbf{k}, -\mathbf{q}) + \hat{\Gamma}(\mathbf{k}, -\mathbf{k}') &= 0. \end{aligned} \quad (\text{D29})$$

The previous equations can be transformed into two uncoupled Fredholm integral equations, whose formal solution is a Liouville-Neumann series.

APPENDIX E: FLUCTUATING HYDRODYNAMICS, BULK EQUATIONS AND STRUCTURE FACTORS

Starting from the bulk fluctuating hydrodynamic equations reported in Eqs. (1) and (2)

$$\partial_t \rho_t + \nabla \cdot (\rho_t \mathbf{v}_b) = 0, \quad \partial_t \rho + \nabla \cdot (\rho \mathbf{v}_b) = \nabla \cdot (\mathcal{D} \nabla \mu + \Psi), \quad (\text{E1})$$

$$\partial_t (\rho_t \mathbf{v}_b) + \nabla \cdot (\rho_t \mathbf{v}_b \mathbf{v}_b) = -\nabla P + \nabla \cdot \{\eta [\nabla \mathbf{v}_b + (\nabla \mathbf{v}_b)^T] + \Sigma\}, \quad (\text{E2})$$

we want to quantify the equilibrium structure factors (static covariances) of the fluctuating fields. These can be obtained by linearizing the above equations around a uniform reference state, $\rho_t = \rho_{t0} + \delta \rho_t$, $C = C_0 + \delta C$, $\mathbf{v}_b = \delta \mathbf{v}_b$, $P = P_0 + \delta P = P_0 + c_s^2 [\delta \rho_t - \rho_{t0} \beta \delta C]$, and then applying a spatial Fourier transform [9]. In the notation used, β is known as the ‘‘solulal expansion’’ coefficient,

$$\rho_{t0} \beta = \left(\frac{\partial \rho_t}{\partial C} \right)_P, \quad (\text{E3})$$

while c_s^2 is the squared speed of sound. Notice that all partial derivatives are evaluated on the uniform reference state. The results for the structure factors are found to be [9, 11]

$$S_{\rho_t, \rho_t}(\mathbf{k}) = \rho_{t0} k_B T \left(\frac{1}{c_s^2} + \frac{\beta^2}{\mu_C} \right), \quad S_{C, C}(\mathbf{k}) = \frac{k_B T}{\rho_{t0} \mu_C}, \quad S_{\rho_t, C}(\mathbf{k}) = \beta \frac{k_B T}{\mu_C}, \quad (\text{E4})$$

where we have indicated with $\mu_C = (\partial\mu/\partial C)_P$. For the model that we consider explicitly in the numerical simulations, the bulk pressure P and the chemical potential μ assume the form [54]

$$P(\rho_t, C) = c_T^2 \rho_t + c_T^2 \mathcal{G} \rho \rho' = c_T^2 \rho_t + c_T^2 \mathcal{G} \rho_t^2 C(1 - C), \quad (\text{E5})$$

$$\mu(\rho_t, C) = c_T^2 \ln \rho - c_T^2 \ln \rho' + c_T^2 \mathcal{G}(\rho' - \rho) = c_T^2 \ln \left(\frac{C}{1 - C} \right) + c_T^2 \mathcal{G} \rho_t (1 - 2C). \quad (\text{E6})$$

A further quantity of interest is

$$\mu_C = \left(\frac{\partial\mu}{\partial C} \right)_P = \frac{c_T^2}{C_0(1 - C_0)} - 2c_T^2 \mathcal{G} \rho_{t0} + c_T^2 \mathcal{G}(1 - 2C_0) \left(\frac{\partial\rho_t}{\partial C} \right)_P. \quad (\text{E7})$$

At constant pressure, $dP = 0$, and we find

$$0 = c_T^2 d\rho_t + 2c_T^2 \mathcal{G} \rho_{t0} C_0(1 - C_0) d\rho_t + c_T^2 \mathcal{G} \rho_{t0}^2 (1 - 2C_0) dC \quad (\text{E8})$$

and hence we can find $(\partial\rho_t/\partial C)_P$, which is defining the parameter β :

$$\rho_{t0} \beta = \left(\frac{\partial\rho_t}{\partial C} \right)_P = - \frac{\mathcal{G} \rho_{t0}^2 (1 - 2C_0)}{1 + 2\mathcal{G} \rho_{t0} C_0 (1 - C_0)}. \quad (\text{E9})$$

Equation (E7) then becomes

$$\mu_C = \frac{c_T^2}{C_0(1 - C_0)} \frac{1 - C_0(1 - C_0) \rho_{t0}^2 \mathcal{G}^2}{1 + 2C_0(1 - C_0) \rho_{t0} \mathcal{G}}. \quad (\text{E10})$$

The square of the sound speed is defined in terms of the bulk pressure

$$c_s^2 = \left(\frac{\partial P}{\partial \rho_t} \right)_C = c_T^2 [1 + 2\mathcal{G} \rho_{t0} C_0 (1 - C_0)]. \quad (\text{E11})$$

Other relations of interest are provided by

$$\frac{1}{c_s^2} + \frac{\beta^2}{\mu_C} = \frac{1}{c_T^2} \frac{1 - 2C_0(1 - C_0) \rho_{t0} \mathcal{G}}{1 - C_0(1 - C_0) \rho_{t0}^2 \mathcal{G}^2}, \quad (\text{E12})$$

$$\frac{\beta}{\mu_C} = - \frac{C_0(1 - C_0)}{c_T^2} \frac{(1 - 2C_0) \rho_{t0} \mathcal{G}}{1 - C_0(1 - C_0) \rho_{t0}^2 \mathcal{G}^2}. \quad (\text{E13})$$

Based on Eqs. (E4) and the results obtained in Eqs. (E10)–(E13), the structure factors are evaluated

$$S_{\rho_t, \rho_t}(\mathbf{k}) = \rho_{t0} k_B T \left(\frac{1}{c_s^2} + \frac{\beta^2}{\mu_C} \right) = \frac{\rho_{t0} k_B T}{c_T^2} \frac{1 - 2C_0(1 - C_0) \rho_{t0} \mathcal{G}}{1 - C_0(1 - C_0) \rho_{t0}^2 \mathcal{G}^2}, \quad (\text{E14})$$

$$S_{C, C}(\mathbf{k}) = \frac{k_B T}{\rho_{t0} \mu_C} = \frac{k_B T C_0(1 - C_0)}{c_T^2 \rho_{t0}} \frac{1 + 2C_0(1 - C_0) \rho_{t0} \mathcal{G}}{1 - C_0(1 - C_0) \rho_{t0}^2 \mathcal{G}^2}, \quad (\text{E15})$$

$$S_{\rho_t, C}(\mathbf{k}) = \beta \frac{k_B T}{\mu_C} = - \frac{k_B T C_0(1 - C_0)}{c_T^2} \frac{(1 - 2C_0) \rho_{t0} \mathcal{G}}{1 - C_0(1 - C_0) \rho_{t0}^2 \mathcal{G}^2}, \quad (\text{E16})$$

which are in agreement with the $\mathbf{k} \rightarrow \mathbf{0}$ limit obtained from Eqs (59). Repeating the calculations including the higher-order derivatives in the forcing terms would lead to a wave-vector-dependent sound speed and chemical potential, and the corresponding linearized hydrodynamic equations would also predict a \mathbf{k} dependency of the structure factors [13]. Alternatively, one could use a free-energy functional [54] made up of an ideal part plus interaction terms (directly related to the forcing terms) and study the density fluctuations around equilibrium [12].

-
- [1] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, UK, 1997).
 [2] J. Lyklema, *Fundamentals of Interface and Colloid Science* (Academic Press, London, 1991).
 [3] W. B. Russel, D. A. Saville, and W. R. Schowalter, *Colloidal Dispersions* (Cambridge University Press, Cambridge, UK, 1995).

- [4] L. E. Reichl, *A Modern Course in Statistical Physics* (Wiley, New York, 1998).
 [5] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon, New York, 1959).
 [6] P. C. Hohenberg and B. I. Halperin, *Rev. Mod. Phys.* **49**, 435 (1977).
 [7] In the viscous stress tensor, $\eta(\nabla\mathbf{v}_b + (\nabla\mathbf{v}_b)^T - \frac{2}{D}\mathbb{1}(\nabla \cdot \mathbf{v}_b)) + \eta_b \mathbb{1}(\nabla \cdot \mathbf{v}_b)$, we have assumed a bulk viscosity η_b such that

- $\frac{2}{D}\eta = \eta_b$. This is realized with the LBE when the relaxation times of shear and bulk modes are the same (Sec. VI).
- [8] M. Sega, M. Sbragaglia, S. S. Kantorovich, and A. O. Ivanov, *Soft Matter* **9**, 10092 (2013).
- [9] J. M. Ortiz de Zárate and J. V. Sengers, *Hydrodynamic Fluctuations in Fluids and Fluid Mixtures* (Elsevier Science, Amsterdam, 2006).
- [10] All these partial derivatives are evaluated on the uniform reference state.
- [11] A. Donev, A. J. Nonaka, Y. Sun, T. Fai, A. L. Garcia, and J. B. Bell, *Commun. Appl. Math. Comput. Sci.* **9**, 47 (2014).
- [12] M. Gross, R. Adhikari, M. E. Cates, and F. Varnik, *Phys. Rev. E* **82**, 056714 (2010).
- [13] M. Gross, M. E. Cates, F. Varnik, and R. Adhikari, *J. Stat. Mech.: Theory. Exp.* (2011) P03030.
- [14] A. J. C. Ladd, *J. Fluid. Mech.* **271**, 285 (1994).
- [15] J. B. Bell, A. L. Garcia, and S. A. Williams, *Phys. Rev. E* **76**, 016708 (2007).
- [16] A. Donev, E. Vanden-Eijnden, A. L. Garcia, and J. B. Bell, *Commun. Appl. Math. Comput. Sci.* **5**, 149 (2010).
- [17] J. B. Bell, A. Garcia, and S. Williams, *ESAIM: Mathematical Modelling and Numerical Analysis* **44**, 1085 (2010).
- [18] S. P. Thampi, I. Pagonabarraga, and R. Adhikari, *Phys. Rev. E* **84**, 046709 (2011).
- [19] G. Gonnella, E. Orlandini, and J. M. Yeomans, *Phys. Rev. E* **59**, R4741 (1999).
- [20] A. Prosperetti and G. Tryggvason, *Computational Methods for Multiphase Flow* (Cambridge University Press, Cambridge, UK, 2007).
- [21] C. Brennen, *Fundamentals of Multiphase Flow* (Cambridge University Press, Cambridge, UK, 2005).
- [22] A. Onuki, *Phase Transition Dynamics* (Cambridge University Press, Cambridge, UK, 2002).
- [23] R. Benzi, S. Succi, and M. Vergassola, *Phys. Rep.* **222**, 145 (1992).
- [24] S. Chen and G. D. Doolen, *Annu. Rev. Fluid Mech.* **30**, 329 (1998).
- [25] J. Zhang, *Microfluidics & Nanofluidics* **10**, 1 (2011).
- [26] C. K. Aidun and J. R. Clausen, *Annu. Rev. Fluid Mech.* **42**, 439 (2010).
- [27] B. B. Kadomtsev, *Sov. Phys. JETP* **5**, 771 (1957).
- [28] M. Bixon and R. Zwanzig, *Phys. Rev.* **187**, 267 (1969).
- [29] R. F. Fox and G. E. Uhlenbeck, *Phys. Fluids* **13**, 1893 (1970).
- [30] R. F. Fox and G. E. Uhlenbeck, *Phys. Fluids* **13**, 2881 (1970).
- [31] J. Logan and M. Kac, *Phys. Rev. A* **13**, 458 (1976).
- [32] A. J. F. Siegert A. J. F., *Phys. Rev.* **76**, 1708 (1949).
- [33] Y. L. Klimontovich, *Sov. Phys. Usp.* **16**, 512 (1974).
- [34] J. W. Dufty and M. H. Ernst, in *Pattern Formation and Lattice Gas Automata (Proceedings Nato Workshop, Waterloo, Canada, June 7–12, 1993)*, edited by A. Lawniczak and R. Kapral, Fields Institute Communications 6 (American Math. Society, Providence, RI, 1993).
- [35] R. Adhikari, K. Stratford, M. E. Cates, and A. J. Wagner, *Europhys. Lett.* **71**, 473 (2005).
- [36] B. Dünweg, U. D. Schiller, and A. J. C. Ladd, *Phys. Rev. E* **76**, 036704 (2007).
- [37] G. Kaehler and A. J. Wagner, *Phys. Rev. E* **87**, 063310 (2013).
- [38] Z. Zelko and B. Dünweg, [arXiv:1402.2920](https://arxiv.org/abs/1402.2920).
- [39] L. Onsager and S. Machlup, *Phys. Rev.* **91**, 1505 (1953).
- [40] S. R. de Groot and P. Mazur, *Non-equilibrium Thermodynamics* (North-Holland, Amsterdam, 1962).
- [41] D. Forster, *Hydrodynamic Fluctuations, Broken Symmetry, and Correlation Functions* (Perseus Books, 1990).
- [42] J.-P. Hansen and I. R. McDonald, *Theory of Simple Liquids*, 3rd ed. (Academic Press, San Diego, 2006).
- [43] X. Shan, X. F. Yuan, and H. Chen, *J. Fluid Mech.* **550**, 413 (2006).
- [44] H. Grad, *Pure Appl. Math.* **2**, 331 (1949).
- [45] H. Grad, *Pure Appl. Math.* **2**, 325 (1949).
- [46] R. Zwanzig, *Non Equilibrium Statistical Mechanics* (Oxford University Press, Oxford, UK, 2001).
- [47] P. Asinari, *Phys. Rev. E* **77**, 056706 (2008).
- [48] M. E. McCracken and J. Abraham, *Phys. Rev. E* **71**, 046704 (2005).
- [49] P. L. Bhatnagar, E. P. Gross, and M. Krook, *Phys. Rev.* **94**, 511 (1954).
- [50] X. Shan and H. Chen, *Phys. Rev. E* **47**, 1815 (1993).
- [51] X. Shan and H. Chen, *Phys. Rev. E* **49**, 2941 (1994).
- [52] M. Sbragaglia and D. Belardinelli, *Phys. Rev. E* **88**, 013306 (2013).
- [53] S. Bastea, R. Esposito, J. L. Lebowitz, and R. Marra, *Phys. Rev. Lett.* **89**, 235701 (2002).
- [54] R. Benzi, M. Sbragaglia, S. Succi, M. Bernaschi, and S. Chibbaro, *J. Chem. Phys.* **131**, 104903 (2009).
- [55] R. W. Nash, R. Adhikari, and M. E. Cates, *Phys. Rev. E* **77**, 026709 (2008).
- [56] From Eq. (31), for a mode T_a of order n in the velocity, the corresponding Φ_a is given in terms of lower-order moments m_a . By induction, if the lower-order m_a are already known in terms of the new moments \bar{m}_a of the same order, then, using (65)–(67), we can express the m_a of order n in terms of the \bar{m}_a of order n . By inserting such expressions in the Φ_a of order $n + 1$ we can then express it in terms of the \bar{m}_a .
- [57] M. Sbragaglia, R. Benzi, L. Biferale, S. Succi, K. Sugiyama, and F. Toschi, *Phys. Rev. E* **75**, 026702 (2007).
- [58] M. Gross, R. Adhikari, M. E. Cates, and F. Varnik, *Philos. Trans. R. Soc., A* **369**, 2274 (2010).
- [59] W. H. Press and S. A. Teukolsky, S. A., W. T. Vetterling and B. P. Flannery, *Numerical Recipes: The Art of Scientific Computing* (Cambridge University Press, New York, 2007).
- [60] M. Grant and R. C. Desai, *Phys. Rev. A* **27**, 2577 (1983).
- [61] Safran, *Statistical Thermodynamics of Surfaces, Interfaces and Membranes* (Westview Press, Boulder, CO, 2003).
- [62] P. E. Kloeden and E. Platen, *Numerical Solution of Stochastic Differential Equations* (Springer, Berlin, 1999).
- [63] A. W. C. Lau and T. C. Lubensky, *Phys. Rev. E* **76**, 011123 (2007).
- [64] B. Dünweg and A. J. C. Ladd, *Adv. Comput. Sim. Soft Mater. Sci.* **III 221**, 89 (2009).
- [65] D. Chiappini, G. Bella, S. Succi, F. Toschi, and S. Ubertini, *Commun. Comput. Phys.* **7**, 423 (2010).
- [66] M. Gross, N. Moradi, G. Zikos, and F. Varnik, *Phys. Rev. E* **83**, 017701 (2011).