

Anomalous scaling in random shell models for passive scalars

A. Wirth¹ and L. Biferale²

¹*Observatoire de Nice, CNRS, Boîte Postale 229, 06304 Nice Cedex 4, France*

²*Dipartimento di Fisica, Università di Roma "Tor Vergata," Via della Ricerca Scientifica 1, 00133 Roma, Italy*

(Received 7 May 1996)

A shell-model version of Kraichnan's [Phys. Rev. Lett. **72**, 1016 (1994)] passive scalar problem is introduced which is inspired by the model of Jensen, Paladin, and Vulpiani [Phys. Rev. A **45**, 7214 (1992)]. As in the original problem, the prescribed random velocity field is Gaussian and δ correlated in time, and has a power-law spectrum $\propto k_m^{-\xi}$, where k_m is the wave number. Deterministic differential equations for second- and fourth-order moments are obtained and then solved numerically. The second-order structure function of the passive scalar has normal scaling, while the fourth-order structure function has anomalous scaling. For $\xi = \frac{2}{3}$ the anomalous scaling exponents ζ_p are determined for structure functions up to $p = 16$ by Monte Carlo simulations of the random shell model, using a stochastic differential equation scheme, validated by comparison with the results obtained for the second- and fourth-order structure functions. [S1063-651X(96)13510-8]

PACS number(s): 47.27.-i

I. INTRODUCTION

A most striking property of a passive scalar quantity advected by a fully developed turbulent flow is its spatiotemporal intermittent behavior (see Ref. [1] for a recent review). The time evolution of the scalar field θ is described by the partial differential equation

$$\partial_t \theta + (\mathbf{v} \cdot \nabla) \theta = \kappa \Delta \theta + f, \quad (1)$$

where \mathbf{v} is the advecting velocity field, κ is the molecular diffusivity, and f is an external forcing.

Intermittency in passive scalars advected by highly chaotic flows is thought to be connected to the existence of self-similar processes transferring fluctuations from large to small scales.

Self-similarity is observed in the power-law behavior of structure functions $\tilde{S}_p(r)$ in the inertial range; that is, moments of θ increments at scales where neither external forcing nor molecular damping are acting

$$\tilde{S}_p(r) \equiv \langle |\theta(x) - \theta(x+r)|^p \rangle \sim r^{\zeta_p}. \quad (2)$$

The set of scaling exponents ζ_p fully characterizes intermittency. In particular, deviations from the dimensional (linear) behavior $\zeta_{2p} = p\zeta_2$ are evidence of a nontrivial scalar transfer among scales, in analogy to the energy intermittent cascade in turbulent flows (see Ref. [2] for a recent review on this subject).

Another important issue is connected to the questioned universal character of scalar fluctuations in the inertial range. Universality should be the consequence of local, or quasilocal, interactions among scales, and it should be reflected in a strong robustness of the scaling exponents ζ_p against variation of forcing and/or dissipation mechanisms. This idea dates back to the phenomenological work of Obukhov and Corrsin [3].

Statistics of the passive scalar must be strongly related to the properties of the advecting velocity fields; therefore, the only realistic case would be to study the "full problem" given by (1) together with the corresponding Navier-Stokes

equations, in the limit of high Reynolds number, which describes the time evolution of a fully developed turbulent velocity field.

Due to the lack of knowledge about the statistical properties of a fully developed turbulent flow the "full problem" cannot yet be attacked. Nevertheless, Eq. (1), with a prescribed velocity field of suitable statistical properties, could be a good "training ground" for understanding general mechanisms (if any) underlying the presence of anomalous scaling in fluid dynamics.

Recently, much attention has been paid in this direction by studying the intermittency properties of a passive scalar advected by a velocity field which is Gaussian, δ -correlated in time, and which has self-similar spatial correlations. In Refs. [4–11] the two-point velocity-field correlation function was set to $\langle v_i(x,t)v_j(x',t') \rangle = \delta(t-t')D_{ij}(x-x')$, with $D(x) = D(0) - \hat{D}(x)$. Here, \hat{D} is the d -dimensional velocity-field structure function

$$\hat{D}_{ij}(x) = D_0 |x|^\xi [(d-1+\xi)\delta_{ij} - \xi x_i x_j |x|^{-2}], \quad (3)$$

where the scaling exponent ξ of the second-order velocity structure function, ξ with $0 < \xi < 2$, is a free parameter. Higher-order velocity-field correlation functions are fixed by the Gaussian assumption.

Such a choice is far from being realistic because of the assumed fast-time decorrelation and Gaussianity. Therefore, no quantitative agreement with the intermittent properties of the "full problem" must be expected. Nevertheless, in Refs. [4–7, 9, 10, 12] many interesting analytical and phenomenological results for this toy model have been obtained, giving the impression that some additional new insight into the "intermittency problem" has been gained (see Ref. [13] for related results in a passive vector case).

The big advantage of this toy model is that, due to the δ correlation in time, all equations for equal-time n -point correlations are formally closed, that is, the n th-order correlation depends only on lower-order correlations. This fact was stressed by Kraichnan [4,14], who gave closed expression for $\tilde{S}_2(r)$. In Ref. [4] a theory for all structure functions was

proposed, and an explicit formula for ζ_p given. The main physical outcome was that all structure functions of order greater than 2 have intermittency corrections, and that intermittency is connected to some nontrivial matching between advective and diffusive properties of the model. Indeed, for structure functions of order greater than 2, it is no longer possible to clearly separate the contributions from diffusion and advection terms in the generation of anomalous scaling.

In Refs. [5, 7, 9] it was shown that intermittency of the scalar structure functions is connected to properties of the null space of the linear operators describing the inertial-scale evolution of multipoint passive scalar moments. Moreover, perturbative expressions of intermittency corrections, as a function of the parameter ξ and as a function of the inverse of the dimensionality of the system, were given [5,7,9]. In both cases careful analysis of matching conditions at infrared (IR) and ultraviolet (UV) scales must be taken into account. Universality in the scaling exponents is still preserved, while constants in front of the power laws acquire dependencies on the forcing and on the passive scalar dissipation. How far all these results can be freely extended to the “full problem” is difficult to say. Certainly, the scaling exponents have a strong quantitative dependence on the statistical properties of the prescribed velocity field. Influence of a short but nonzero correlation time was studied in Ref. [8] and found to be important, while from a phenomenological point of view it is difficult to say if the perturbation expansion of Ref. [5] can be extended up to the value $\xi = \frac{2}{3}$ which mimics the spatial energy spectrum of true turbulence. (Let us notice that, even for $\xi = \frac{2}{3}$, the assumed δ correlation in time for the velocity makes the classical Obukhov-Corrsin theory inapplicable to the passive scalar two-point correlation function [14].)

An important drawback of the Gaussian δ -correlated toy model is that numerical simulations seem even more difficult than a direct numerical simulation of the “full problem.” This is because (1) is a stochastic differential equations (SDE) the solution of which requires huge computer resources, because of the massive use of random number generators and also because of the paramount difficulties in the implementation of higher than first-order discretization schemes [15]. The state of the art of SDE numerical simulations is hardly compatible with the high spatial resolution needed to study scaling properties of the solution of (1) with a δ -correlated in-time velocity field (see, however, Ref. [6] for an attempt at overcoming these difficulties, albeit at the price of introducing some anisotropy into the velocity field).

In this spirit, we have tried to work with the simplest toy model which has some connection to the physics of the “full problem,” while at the same time aiming for something more tractable analytically and/or numerically. We have thus investigated the intermittency properties of a shell model for a passive scalar advected by a prescribed stochastic velocity field. Shell models (see Ref. [16] for a pedagogical introduction) have already been successful in helping to understand many issues connected to fully developed turbulence [17–22].

The problem of defining a shell model for the advection of a passive scalar by a deterministic and chaotic velocity field has been already investigated in Ref. [23]. Here we are

going to present our numerical and analytical results obtained in a weakly modified version of the shell model defined in Ref. [23], but using a δ -correlated surrogate for the advecting velocity field. The main drawback of our model is that, following shell-model philosophy, it retains only nearest- and next-nearest-neighbor interactions in the shell index. Having only semilocal interaction in Fourier space could be a problem for describing the physics of passive fluctuations for those types of velocity-field correlations which induce strong nonlocal effects. For example, simple dimensional considerations tell us that the limit of very small ξ should be dominated by strong nonlocal effects (which leads to an ultraviolet divergence of the eddy diffusivity in the limit $\xi \rightarrow 0$ [5]). Therefore we do not expect that our shell model could properly mimic the physics of the partial differential equation (PDE) (1) in the range of parameters choice $0 < \xi \ll 1$. We are quite confident that our model captures very well the passive scalar physics for $\xi = O(1)$: that is, when dominant interactions become more local in Fourier space, as is the case for true turbulence [24]. The major advantage of using a shell model is that, now, reliable numerical simulations become feasible, allowing nonperturbative determinations of scaling exponents up to high orders (in this paper we present results up to order 16). Furthermore, having reliable numerical simulations should open the possibility of relaxing some of the constraints put on the statistical properties of the velocity field, thereby allowing investigation of increasingly realistic problems.

We found that our shell model shares many properties with the original SDE (1). Among them, the most important are that (i) the second-order structure function has normal scaling, (ii) all structure functions of orders larger than 2 have anomalous corrections, and (iii) there is a remarkable coupling between both UV and IR scales with inertial terms in the closed equation satisfied by fourth-order shell correlations. This result suggests that the anomalous scaling is due to nontrivial contributions of integral and diffusive scales on the inertial-range properties. Moreover, we find that anomalous behavior tends to vanish when approaching the laminar regime, $\xi = 2$, for the advecting velocity field.

Even though we have not obtained an analytical expression for the anomalous scaling exponents, we are confident that many useful insights can be obtained by further investigation of the present model, or an even simpler one. For example, it should be possible to define some exactly solvable models, where the shell-velocity correlations are chosen such as to give exactly solvable (but nontrivial) linear operators describing the structure functions-inertial properties.

The paper is organized as follows: in Sec. II we introduce the random passive scalar shell model, and we discuss some problems connected with its stochastic differential equation (SDE) formulation. In Sec. III we explicitly derive the closed deterministic equations for second- and fourth-order moments, which are numerically solved in Sec. IV to obtain the scaling exponents of second- and fourth-order structure functions. Monte Carlo simulations of the SDE are presented in Sec. V. Concluding remarks and suggestions for further work are given in Sec. VI.

II. THE MODEL AND ITS STOCHASTIC DIFFERENTIAL EQUATION FORMULATION

We recall the main features of the passive scalar shell model of Ref. [23]. The model is defined in terms of a shell

discretization of the Fourier space in a set of wave numbers defined on a geometric progression $k_n = k_0 2^n$. Passive increments at scale $r_n = k_n^{-1}$ are described by a complex variable $\theta_n(t)$. The time evolution is obtained according to the following criteria: (i) the linear term is a purely diffusive term given by $-\kappa k_n^2 \theta_n$; (ii) the advection term is a combination of the form $k_n \theta_{n'}, u_{n''}$; (iii) interacting shells are restricted to nearest and next-nearest neighbors of n ; and (iv) in the absence of forcing and damping the model conserves the volume in the phase space and the passive energy $E = \sum_n |\theta_n|^2$. Properties (i), (ii), and (iv) are also valid for the original equation (1) in Fourier space, while property (iii) is an assumption of locality of interactions among modes, which is rather well founded as long as the power-law spectra of the passive scalar $E(k) \sim k^{-\alpha}$ has $1 < \alpha < 3$.

Our passive scalar shell model, inspired by Jensen, Paladin, and Vulpiani's model [23], is defined by the following equations ($m = 1, 2, \dots$):

$$\begin{aligned} \left[\frac{d}{dt} + \kappa k_m^2 \right] \theta_m(t) = & i \{ a_m [\theta_{m+1}^*(t) u_{m-1}^*(t) \\ & - \theta_{m-1}^*(t) u_{m+1}^*(t)] \\ & + b_m [\theta_{m-1}^*(t) u_{m-2}^*(t) \\ & + \theta_{m-2}^*(t) u_{m-1}(t)] \\ & + c_m [\theta_{m+2}^*(t) u_{m+1}(t) \\ & + \theta_{m+1}^*(t) u_{m+2}^*(t)] \} + \delta_{m1} f(t), \end{aligned} \quad (4)$$

where the asterisk denotes complex conjugation and

$$k_m = 2^m, \quad a_m = \frac{k_m}{2}, \quad b_m = -\frac{k_{m-1}}{2}, \quad c_m = \frac{k_{m+1}}{2}, \quad (5)$$

with $u_{-1} = u_0 = \theta_{-1} = \theta_0 = 0$ as boundary conditions. The forcing term $\delta_{1m} f(t)$ acts only on the first shell. Note that Eq. (5) is one of possible choices for the parameters a_m , b_m , and c_m , ensuring the conservation of passive scalar energy $\sum_m \theta_m \theta_m^*$ and phase-space volume. In numerical implementations, the model is truncated to a finite number of shells N (here $N=19$) with the additional boundary conditions $u_{N+1} = u_{N+2} = \theta_{N+1} = \theta_{N+2} = 0$.

Our model differs from that of Ref. [23] by the absence of complex conjugation on two of the u factors in the right-hand side of (4). The reason for this change will become clear below. Furthermore, in Ref. [23] the passive scalar model (4) was coupled to the Gledzer-Ohkitani-Yamada (GOY) shell model [17–19] for the nonlinear dynamics of the velocity variables. The GOY model displays multifractal behavior for the u_n 's, and an interesting intermittent behavior is also found for the passive scalar.

Our goal, as in Kraichnan's work [4], is to use a *nonintermittent* velocity field and then to find if the passive scalar is nevertheless intermittent. For this, we assume that the velocity variables $u_m(l)$ and the forcing term $f(t)$ are independent complex Gaussian and δ correlated and in time. Furthermore, as in Ref. [14], we make a scaling assumption for the spectrum of the $u_m(t)$'s, namely

$$\langle u_m(t) u_m^*(t') \rangle = \delta(t-t') D_m, \quad (6)$$

where

$$D_m = k_m^{-\xi} \quad (7)$$

will be called the velocity spectrum. For $f(t)$ we assume $\langle f(t) f(t') \rangle = \delta(t-t')$.

As long as the velocity variables $u_m(t)$ have a finite correlation time, and hence, smooth sample paths, there is no particular difficulty in giving a meaning to the set (4) of random ordinary differential equations (ODE's). A well-known difficulty arises with ODE's having δ -correlated in-time coefficients: the mathematical meaning of the equation is ambiguous [15]. The physicist's viewpoint, and thus our own, is to define the solution of such "stochastic differential equations" to be the limit, as the correlation time tends to zero, of the solution of a random ODE with nonwhite (colored) coefficients. In the mathematical literature on stochastic differential equations this is called the Stratonovich version of stochastic calculus [15]. When numerically solving a SDE such as (4), one cannot use standard numerical methods which assume *smooth* coefficients. It would also be highly impractical to use a smooth approximation of the coefficients with a small correlation time, since this requires time steps much smaller than the correlation time. Fortunately, there is an alternative formulation of the SDE, the Ito version, which overcomes this difficulty. The solutions of SDE's are Markov diffusion process which can be characterized by their diffusion and drift coefficients (also called Fokker-Planck coefficients). The so-called Ito equation [15] encodes this information in a way which leads naturally to efficient numerical schemes.

Let us briefly explain how this Ito equation is obtained. For this it is better to work with an abstract form of the starting equation (with the forcing term omitted):

$$\frac{d\theta}{dt} = \mathbf{M}_0 \theta + \frac{1}{\epsilon} \mathbf{M}_1 \left(\frac{t}{\epsilon^2} \right) \theta. \quad (8)$$

Here, \mathbf{M}_0 is a deterministic operator and \mathbf{M}_1 a random Gaussian operator with a finite correlation time. The ϵ scaling is chosen in such away that, for $\epsilon \rightarrow 0$, the \mathbf{M}_1 operator becomes δ correlated in time. Just as ordinary white noise may be written as the derivative, in the sense of distributions, of the Brownian motion process, we can write

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \mathbf{M}_1 \left(\frac{t}{\epsilon^2} \right) = \frac{d\mathbf{W}}{dt}, \quad (9)$$

where $\mathbf{W}(t)$ is an operator-valued Brownian motion. The Ito SDE associated with (8) is

$$d\theta = (\mathbf{M}_0 + \mathcal{D}) \theta dt + d\mathbf{W} \theta, \quad (10)$$

where

$$\mathcal{D} = \int_0^\infty \langle \mathbf{M}_1(s) \mathbf{M}_1(0) \rangle ds \quad (11)$$

is called the drift operator. The presence of the drift term in Eq. (10) is easily understood: a Neumann expansion of $\theta(t$

$+dt) - \theta(t)$ in powers of \mathbf{M}_1 , starting from Eq. (8), gives a second-order term, which is $O(dt^2)$ for fixed ϵ but becomes $D\theta dt$ when the limit $\epsilon \rightarrow 0$ is taken. The drift term may also be obtained by using the Ito formula [15] or Gaussian integration by parts (see, e.g., Ref. [2], Sec. 4.1). The term $d\mathbf{W}\theta$ is called the diffusion term. Since Brownian motion has independent increments over nonoverlapping intervals, the operator $d\mathbf{W}$, which is $O(\sqrt{dt})$ and has zero mean, is indepen-

dent of the past and, thus, of θ . It follows that the diffusion term has vanishing mean. Hence, the mean of θ satisfies

$$\frac{d\langle\theta\rangle}{dt} = (\mathbf{M}_0 + \mathcal{D})\langle\theta\rangle. \quad (12)$$

The Ito SDE associated with (4) reads

$$\begin{aligned} d\theta_m(t) = & [(a_m b_{m+1} + b_m c_{m-2})D_{m-1} + (-a_m c_{m-1} + b_{m+2} c_m)D_{m+1} + a_{m-1} b_m D_{m-2} - a_{m+1} c_m D_{m+2} - \kappa k_m^2] \theta_m(t) dt \\ & + i \{ a_m [\theta_{m+1}^*(t) \alpha_{m-1} dW_{m-1}^*(t) - \theta_{m-1}^*(t) \alpha_{m+1} dW_{m+1}^*(t)] + b_m [\theta_{m-1}^*(t) \alpha_{m-2} dW_{m-2}^*(t) \\ & + \theta_{m-2}^*(t) \alpha_{m-1} dW_{m-1}^*(t)] + c_m [\theta_{m+2}^*(t) \alpha_{m+1} dW_{m+1}^*(t) + \theta_{m+1}^*(t) \alpha_{m+2} dW_{m+2}^*(t)] \} + \delta_{m,1} dW_f(t). \end{aligned} \quad (13)$$

Here $\alpha_m = k_m^{\xi/2}$ and $D_m = \alpha_m^2/2$, and the $W_m(t)$'s and $W_f(t)$ are independent identically distributed complex-valued Brownian motion functions, normalized in such a way that $\langle |W_m(t)|^2 \rangle = \langle |W_f(t)|^2 \rangle = t$. It is noteworthy that the drift term in Eq. (13), the top two lines on the right-hand side, which involves the sum of the energies in the neighboring and next-neighboring shells, may be viewed as an eddy-diffusivity term. For Kraichnan's original equation (1), the Ito equation approach has a scale-independent eddy diffusivity [26]. Here, it is proportional to $k_n^{-\xi}$. The difference stems from the absence, in the shell model, of direct interactions between widely separated scales. We also observe that, if we had used the original model of Ref. [23] with complex conjugates on all the u factors, the drift operator would involve nondiagonal elements coupling different shells, a situation we avoided. Indeed, the drift operator is the mean value of all double stochastic integrals of the stochastic Taylor expansion [15]. By Taylor expanding (13) and using $E(\int_0^t dW dW) = 0$ and $E(\int_0^t dW^* dW) = t/2$, the nondiagonality of the drift operator in Ref. [23] is easily verified.

III. EQUATIONS FOR THE SECOND- AND FOURTH-ORDER MOMENTS

In this and the following sections we are interested in the scaling behavior of the p th-order structure functions:

$$\langle (\theta_m \theta_m^*)^{p/2} \rangle \propto k_m^{-\zeta_p}, \quad (14)$$

where ζ_p is called the scaling exponent of order p . If $\zeta_{2p} = p\zeta_2$, the structure functions are said to have a normal scaling. If $\rho_{2p} = p\zeta_2 - \zeta_{2p} \neq 0$, the scaling of the structure function of order $2p$ is said to be anomalous.

It is well known that from a linear stochastic differential equation with white-noise coefficients, it is possible to obtain exact equations for moments of arbitrary order [14,6]. For example, from the abstract equation (8) one derives the closed equation for the first-order moment (12). Higher-order quantities such as $\theta \otimes \theta, \theta \otimes \theta \otimes \theta, \dots$ also satisfy linear stochastic differential equations with white-noise coefficients,

from which closed equations can be obtained for $\langle \theta \otimes \theta \rangle, \langle \theta \otimes \theta \otimes \theta \rangle$, etc. Such equations may also be obtained by use of Ito calculus [15].

In the shell-model context the moment equations become excessively cumbersome beyond order 4. We have obtained closed equations for

$$E_m \equiv \langle \theta_m \theta_m^* \rangle \quad \text{and} \quad P_{lm} = \langle \theta_l \theta_l^* \theta_m \theta_m^* \rangle. \quad (15)$$

The general structure of these equations is as follows

$$\dot{E}_l = (-2\kappa k_m^2 \delta_{l,m} + A_{l,m}) E_m + F_l, \quad (16)$$

$$\dot{P}_{lm} = [-2\kappa(k_l^2 + k_m^2) \delta_{l,n} \delta_{m,j} + B_{lm,nj}] P_{nj} + G_{lm}. \quad (17)$$

The equations are written in explicit form in Appendix B. Here we stress a few important properties. The forcing F_m is restricted only to the first shell $m=1$. The matrix $A_{l,m}$ is a symmetric band diagonal with a bandwidth 5, and has the following scaling law:

$$A_{l+s,m+s} = k_s^{2-\xi} A_{l,m}. \quad (18)$$

Similarly,

$$B_{l+s;m+s,n+s,j+s} = k_s^{2-\xi} B_{lm,nj}. \quad (19)$$

Straightforward scaling arguments indicate that Eqs. (16) and (17) may possess steady-state solutions with *normal* scaling. Such solutions have

$$E_m \propto k_m^{2-\xi} = k_m^{-\zeta_2} \quad (20)$$

and

$$P_{mm} \propto k_m^{-\zeta_4}, \quad (21)$$

with $\zeta_4 = 2\zeta_2$. In order to find what kind of scaling actually holds, we now resort to numerical solutions of the moment equations (16) and (17). Since these are *deterministic* equations, they can be solved with high accuracy at relatively low cost. This is not the case of the Monte Carlo strategy of Sec. V which allows us, however, to tackle structure functions of high orders.




FIG. 1. Anomalous part ρ_4 of the fourth-order structure function exponent vs velocity scaling exponent ξ with error bars.

IV. CALCULATION OF SECOND- AND FOURTH-ORDER STRUCTURE FUNCTIONS FROM THE MOMENT EQUATIONS

For the numerical solution of Eqs. (16) and (17), we used a c19-shell truncation with values of κ chosen in such a way that the diffusive cutoff is well within the available range of shells. An exponential cutoff was put on the velocity spectrum for shells in the dissipation range. Without cutoff in the velocity spectrum the passive-scalar spectrum has an algebraic tail [25]. The solutions were obtained by time marching until the steady state is reached. Such a steady state is necessarily stable. The scaling parameter ξ was varied in the range $0.2 \leq \xi \leq 2$.

We found that the second-order structure function E_m always displays normal scaling with

$$\zeta_2 = 2 - \xi, \quad (22)$$

the same value as obtained by Kraichnan [6]. To investigate the nature of the scaling of the fourth-order structure function P_{mmm} ; we plotted it against the second-order structure function, following the now standard, extended self-similarity (ESS) procedure [26]. This gave us the anomalous part $\rho_4 = 2\zeta_2 - \zeta_4$ of the scaling exponent, plotted in Fig. 1 against ξ . It is clearly seen that the scaling is anomalous ($\rho_4 > 0$), the anomaly is a decreasing function of ξ and disappears as $\xi \rightarrow 2$, the ‘‘laminar’’ limit as in Ref. [4].

For small values of ξ , interactions become more and more nonlocal, and a higher number of shells would be needed to have a sufficiently large inertial range. [The number of shells in the ‘‘intermediate inertial-dissipation range,’’ in which the




FIG. 2. Second-order (continuous line) and fourth-order (dashed line) structure functions of the passive scalar calculated from the moment equations, compared to direct numerical Monte Carlo simulations (squares and circles).

local Reynolds number is $O(1)$, grows like $1/\xi$.] Even though our shell model has only local interactions, purely kinematic effects introduce important long-range diffusive corrections in the limit $\xi \rightarrow 0$. This is why we restricted our calculations of ρ_4 for ξ ranging from 0.2 to 2.0.

The fact that normal scaling is obtained for the second-order structure function is not very surprising, this being exactly the same situation as for Kraichnan’s original problem. It is easily checked that the operator A appearing in Eq. (16) has equipartition solutions (E_m is independent of m) in its null space, both for the full operator and its N -shell truncation. Such equipartition solutions have no associated passive-scalar energy flux, and cannot bring about anomalous scaling.

Let us now consider the anomalous scaling for the fourth-order structure functions which are connected to the properties of the operator $B \equiv B_{lm,nj}$. The operator B has again equipartition solutions in its null space, which cannot cause anomalous scaling. We have checked numerically that finite-shell truncations of B have no other eigenvectors in their null space. This only superficially contradicts the interpretation of anomalous scaling as arising from the zero modes of the inertial operator describing the evolution of moments [9]. In our case there are no anomalous zero modes of the operator B , but still we see numerically a very clean intermittent behavior. Where does the observed anomaly come from? One can imagine two interpretations. The first one rests on the physical observation that the operator B is naturally long range: it mixes inertial scales with cutoffs at both the UV and IR ends. Therefore, its inverse involves a nontrivial mixture of contributions from very different scales. This mixture could be the cause of anomalous scaling, thereby defeating




FIG. 3. ESS plot of the p th-order structure function vs second-order structure function ($p=4, 6,$ and 8) for $\xi=\frac{2}{3}$. Only inertial-range shells are shown. Lines are the best linear fits. Statistical errors are of the order of symbol sizes.

naive (local) dimensional analysis. The second interpretation is to imagine that quazero modes (which would become true zero modes in the limit of an infinite inertial range) already dominate the inverse of B . The two interpretations are not in contradiction. Indeed, our truncated system, being always influenced by UV and IR cutoffs, naturally takes into account boundary conditions and therefore never shows true zero modes. A similar scenario takes place in Ref. [5], where zero modes in the infinite-dimensional function space have to be matched with IR and UV physical cutoffs. Let us finally remark that the analysis of the relevant eigenvectors in the eigenspace of the operator B is highly complicated due to the fact that we are looking at the cone of positive functions, which is not a linear space.

V. MONTE CARLO SIMULATIONS FOR STRUCTURE FUNCTIONS

As we noted, the moment-equation strategy becomes impractical for determining structure functions beyond the fourth order. We therefore resort to Monte Carlo simulations of the stochastic shell model in its Ito version (13). We used the “weak-order-one Euler” scheme, the details of which may be found in Appendix A. Roughly, this means interpreting the Ito equation, (13) as a time-difference equation. This scheme is of order 1 (in the time step Δt) for averaged quantities such as the structure functions. Averages are calculated as time averages, assuming ergodicity (we checked that changes in the seed of the random generator do not affect the results). Integrating over a large number of realizations is thus equivalent to integrating over many large-eddy turnover times.




FIG. 4. Scaling exponents ζ_p normalized by ζ_2 for structure functions up to order $p=16$ (circles) for $\xi=\frac{2}{3}$. The straight line represents normal scaling.

First we validated our Monte Carlo simulation by comparison with the results for the second- and fourth-order structure functions obtained from the moment equations in the previous section. The comparison can be seen in Fig. 2. The agreement of the Monte Carlo and moment-based calculations is comparable to the machine precision (single precision). We mention that, in order to maximize the extent of the inertial range and avoid an inertial-diffusive range with algebraic falloff of the passive scalar spectrum [26], we assumed an exponential cutoff on the velocity spectrum.

In all calculations 19 shells were used, the molecular diffusivity was varied between 2^{-10} and 2^{-8} and the time step between 2^{-29} and 2^{-25} . In order to give an example of the quality of the scaling, in Fig. 3 we show the log-log plot of the fourth, sixth, and eighth structure functions versus the second-order structure function. This is done by ESS in order to improve the scaling.

Figure 4 shows the scaling exponents ζ_p determined by least-square fits using ESS up to order $p=16$ for $\xi=\frac{2}{3}$. The error bars are obtained from the least-square fits. In all the calculations the inertial range included eight shells or more. When halving the number of samples in our statistics, we found that the values obtained remained well within the error bars. The graph of ζ_p appears to be linear at values of p beyond 8, with a slope of about 0.29. We do not rule out that the asymptotic linear trend is an artifact due to insufficient statistics. We observe that the ζ_p 's obtained with our Gaussian δ -correlated in-time velocity are less anomalous than those reported in Ref. [23], where the velocity was already multifractal.

VI. CONCLUSIONS

We proposed and studied a shell model for a randomly advected passive scalar. We suggest that anomalous scaling

of structure functions of order greater than 2 is connected to nontrivial dependence of inertial-range properties on both integral and diffusive scales. This is shown, for example, in the closed equation for the fourth-order moments $\langle \theta_m \theta_m^* \theta_l \theta_l^* \rangle$, where inertial-range shell correlations depend explicitly on correlations between distant shells with $m \ll l$. The main advantage of our model is that reliable numerical simulations become feasible. We numerically estimated anomalous exponents for structure functions up to order 16, and for various scaling exponents ξ of the velocity field ($0.2 \leq \xi \leq 2$). Comparing our numerical results with those obtained in a similar passive shell model [23], but advected by a multifractal velocity field, we find large quantitative differences. This is due to the obvious fact that the θ statistics are strongly correlated to the statistics of the advecting field. Nevertheless, the possibility of writing down closed equations for correlations of any order could help in the understanding of intermittency in a more general cases. Unfortunately, we do not see how to implement in our model the kind of perturbation expansion done in Refs. [5, 9]. The main difficulty is the lack of long-range interactions in our model which forbids a proper definition of an eddy diffusivity in the limit $\xi \rightarrow 0$. Some long-range shell models should be introduced and studied if one wants to follow this path. On the other hand, our result for the anomalous correction of fourth-order structure function becomes more and more nonperturbative in the limit $\xi \rightarrow 0$, in qualitative agreement with the Kraichnan theory for passive scalars [4].

Another interesting problem is to find the simplest passive-scalar shell model which has inertial-range intermit-

tency. For example, it is easy to extract from Eq. (4) an even simpler shell model, by retaining only terms with $\theta_{m-2}^*(t)u_{m-1}(t)$ and $\theta_{m+2}^*(t)u_{m+1}(t)$ in the advecting part. This model has the same properties as listed in Sec. I, and some preliminary numerical results indicate that it has similar intermittency corrections.

Let us finally remark that, up to now, all existing work on anomalies driven by a stochastic velocity field has been rooted in some kind of analysis performed in the *physical space*. In contrast, our shell model exists only in a kind of *Fourier space*. Understanding the anomalous scaling in our shell model at a phenomenological level could help in devising a phenomenology of intermittency. For example, ideas connected to the popular inertial-range cascade picture could be usefully revisited.

ACKNOWLEDGMENTS

We would like to express our gratitude to M. Vergassola for many stimulating and interesting discussions. We have also benefited from extensive discussions with R. Benzi, U. Frisch, and A. Noullez. This work was supported by the French Ministère de la Recherche et de la Technologie, by the European Union (Human Capital and Mobility ERBCHRXCT920001), and by the GDR Mécanique des Fluides Numérique. L.B. would like to acknowledge partial support by Ministère de l'Enseignement Supérieur et de la Recherche (France), and the Observatoire de la Côte d'Azur where this work was completed.

APPENDIX A: EULER SCHEME

The weak-order-one Euler scheme [15] associated with the passive-scalar shell model in its Ito form (4) reads

$$\begin{aligned} \theta_m^{n+1} = & \{1 + [(a_m b_{m+1} + b_m c_{m-2})D_{m-1}\Delta t + (-a_m c_{m-1} + b_{m+2} c_m)D_{m+1}\Delta t + a_{m-1} b_m D_{m-2}\Delta t \\ & - a_{m+1} c_m D_{m+2}\Delta t - \kappa k_m^2 \Delta t]\} \theta_m^n + i[a_m(\theta_{m+1}^{n*} \alpha_{m-1} \Delta W_{m-1}^{n*} - \theta_{m-1}^{n*} \alpha_{m+1} \Delta W_{m+1}^{n*}) \\ & + b_m(\theta_{m-1}^{n*} \alpha_{m-2} \Delta W_{m-2}^{n*} + \theta_{m-2}^{n*} \alpha_{m-1} \Delta W_{m-1}^{n*}) + c_m(\theta_{m+2}^{n*} \alpha_{m+1} \Delta W_{m+1}^{n*} + \theta_{m+1}^{n*} \alpha_{m+2} \Delta W_{m+2}^{n*})] + f \delta_{m,1} \Delta W_f^n. \end{aligned} \quad (\text{A1})$$

Here

$$\Delta W_m^n = \sqrt{\Delta t} \eta_m^n, \quad (\text{A2})$$

where the η_m^n are independent identically distributed complex random variables of the form $a + ib$, where a and b are independent Bernoulli variables with values $\pm 1/\sqrt{2}$. The Bernoulli variables are numerically generated by a linear feedback shift register random number generator (see, e.g., [27]).

The choice of Bernoulli variables rather than Gaussian variables is particularly convenient for numerical purposes. It ensures that averaged quantities such as moments are correct to first order in Δt , a choice consistent with the scheme.

APPENDIX B: MOMENT EQUATIONS

Hereafter, we give the detailed form of the equation for the second-order moments $E_n \equiv \langle 0_m 0_m^* \rangle$ and the fourth-order moment $P_{lm} \equiv \langle \theta_l \theta_l^* \theta_m \theta_m^* \rangle$:

$$\dot{E}_m = +(-2\kappa k_m^2 + A_{m,m})E_m + A_{m,m-2}E_{m-2} + A_{m,m-1}E_{m-1} - A_{m,m+1}E_{m+1} + a_{m,m+2}E_{m+2} + F_m, \quad (\text{B1})$$

$$\begin{aligned}
\dot{P}_{l,m} = & -[2\kappa(k_m^2 + k_l^2) - A_{m,m} - A_{l,l} + (\delta_{l-1,m} + \delta_{l-2,m} + \delta_{l+1,m} + \delta_{l+2,m} + \delta_{l,m-1} + \delta_{l,m-2} + \delta_{l,m+1} + \delta_{l,m+2})A_{l,m}]P_{l,m} \\
& + A_{m,m-2}P_{l,m-2} + A_{m,m-1}P_{l,m-1} + A_{m,m+1}P_{l,m+1} + A_{m,m+2}P_{l,m+2} + A_{l,l-2}P_{l-2,m} + A_{l,l-1}P_{l-1,m} + A_{l,l+1}P_{l+1,m} \\
& + A_{l,l+2}P_{l+2,m} + \delta_{l,m}2(A_{m,m-2}P_{m,m-2} + A_{m,m-1}P_{m,m-1} + A_{m,m+1}P_{m,m+1} + A_{m,m+2}P_{m,m+2}) \\
& + (\delta_{1,m}E_m + \delta_{1,l}E_l + \delta_{1,m}\delta_{1,l}E_1)F_1.
\end{aligned} \tag{B2}$$

The following notation has been used:

$$F_1 = \frac{1}{2}, \tag{B3}$$

$$F_l = 0 \forall l \neq 1, \tag{B4}$$

$$A_{m,m-2} = 2b_m^2 D_{m-1}, \tag{B5}$$

$$A_{m,m-1} = 2(a_m^2 D_{m+1} + b_m^2 D_{m-2}), \tag{B6}$$

$$A_{m,m+1} = 2(a_m^2 D_{m-1} + c_m^2 D_{m+2}), \tag{B7}$$

$$A_{m,m+2} = 2c_m^2 D_{m+1}, \tag{B8}$$

$$A_{m,m} = -A_{m+2,m} - A_{m+1,m} - A_{m-1,m} - A_{m-2,m}. \tag{B9}$$

-
- [1] K. R. Sreenivasan, Proc. R. Soc. London Ser. A **434**, 165 (1991).
- [2] U. Frisch, *Turbulence: the Legacy of A. N. Kolmogorov* (Cambridge University Press, Cambridge, 1995).
- [3] A. M. Obukhov, Izr. Akad. SSSR Serv. Geogr. Geofiz. **13**, 58 (1949); S. Corrsin, J. Appl. Phys. **22**, 469 (1951).
- [4] R. H. Kraichnan, Phys. Rev. Lett. **72**, 1016 (1994).
- [5] K. Gawedzki and A. Kupiainen, Phys. Rev. Lett. **75**, 3834 (1995).
- [6] R. H. Kraichnan, V. Yakhot, and S. Chen, Phys. Rev. Lett. **75**, 240 (1995).
- [7] M. Chertkov, G. Falkovich, I. Kolokolov and V. Lebedev, Phys. Rev. E **52**, 4924 (1995); M. Chertkov and G. Falkovich, Phys. Rev. Lett. **76**, 2706 (1996).
- [8] M. Chertkov, G. Falkovich, and V. Lebedev, Phys. Rev. Lett. **76**, 3707 (1996).
- [9] D. Bernard, K. Gawedzki, and A. Kupiainen, Phys. Rev. E **54**, 2564 (1996).
- [10] A. Pumir, Europhys. Lett. **34**, 25 (1996).
- [11] G. L. Eyink (unpublished)
- [12] A. L. Fairhall, O. Gat, V. L'vov, and I. Procaccia, Phys. Rev. E **53**, 3518 (1996).
- [13] M. Vergassola, Phys. Rev. E **53**, R3021 (1996).
- [14] R. H. Kraichnan, Phys. Fluids **11**, 945 (1968).
- [15] P. E. Kloeden and E. Platen, *Numerical Solution of Stochastic Differential Equations* (Springer-Verlag, Berlin, 1992).
- [16] L. Kadanoff, Phys. Today **48** (9), 11 (1995).
- [17] E. B. Gledzer, Dok. Akad. Nauk. SSSR **209**, 1046 (1973) [Sov. Phys. Dokl. **18**, 216 (1973)].
- [18] M. Yamada and K. Ohkitani, J. Phys. Soc. Jpn. **56**, 4210 (1987); Prog. Theor. Phys. **79**, 1265 (1988).
- [19] M. H. Jensen, G. Paladin, and A. Vulpiani, Phys. Rev. A **43**, 798 (1991); D. Pisarenko, L. Biferale, D. Courvoisier, U. Frisch, and M. Vergassola, Phys. Fluids A **5**, 2533 (1993).
- [20] R. Benzi, L. Biferale, and G. Parisi, Physica D **65**, 163 (1993).
- [21] L. Kadanoff, D. Lohse, J. Wang, and R. Benzi, Phys. Fluids **7**, 617 (1995).
- [22] R. Benzi, L. Biferale, R. Kerr, and E. Trovatore, Phys. Rev. E **53**, 3541 (1996).
- [23] M. H. Jensen, G. Paladin, and A. Vulpiani, Phys. Rev. A **45**, 7214 (1992).
- [24] H. A. Rose and P. Sulem, J. Phys. (Paris) **39**, 441 (1978).
- [25] U. Frisch and A. Wirth, Europhys. Lett. (to be published).
- [26] R. Benzi, S. Ciliberto, R. Tripiccion, C. Baudet, F. Massaioli, and S. Succi, Phys. Rev. E **48**, R29 (1993).
- [27] D. E. Knuth, *Seminumerical Algorithms* (Addison-Wesley, Reading, MA, 1981).